

# CURRENT ISSUES IN NEUROTOXICOLOGY

## *Editors*

**Antonio Mutti**

*Institute of Clinical Medicine and Nephrology  
University of Parma  
Via A. Gramsci 14, 43100 Parma, Italy*

**Luigi Manzo**

*Department of Internal Medicine  
Toxicology Division  
University of Pavia Medical School  
Pavia, Italy*

**Lucio G. Costa**

*Department of Environmental Health  
University of Washington  
SC-34, Seattle, Washington 98195 USA*

**Joan M. Cranmer**

*Department of Pediatrics - ACH #512  
University of Arkansas for Medical Sciences  
800 Marshall Street  
Little Rock, Arkansas 72202-3591 USA*

*Proceedings of the 3rd Meeting of the International Neurotoxicology Association  
held July 1-5, 1991 in Salsomaggiore Terme (Parma), Italy*

**INTOX PRESS, INC., Publishers**

COPYRIGHT © 1992 BY INTOX PRESS, INC.  
ISBN: 0-911369-16-3

Printed in the United States of America on acid free paper

INTOX PRESS, INC.  
Little Rock, Arkansas USA

All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, microfilming, recording, or any information storage and retrieval system, without permission in writing from the Publisher. Direct all inquiries to Intox Press, Inc., P. O. Box 24865, Little Rock, Arkansas 72221 Telephone: (501) 227-8622; FAX: (501) 224-1947.

---

*Library of Congress Cataloging-in-Publication Data*

(Books in the neurotoxicology series; v. 18)

Current issues in neurotoxicology.

Includes bibliographies and index.

1. Neurotoxicology. I. Mutti, Antonio. II. Costa, Lucio G.  
III. Manzo, Luigi. IV. Cranmer, Joan M. V. Title. (DNLM:  
1. Neurotoxicology. 2. Toxicology. 3. Nervous system.  
VI. Series. Books in the neurotoxicology series.  
ISBN: 0-911369-16-3. LC 83-080660

---

Printing: 1 2 3 4 5 6 7 8 9

# CONTENTS

## I. FOREWORD AND ACKNOWLEDGEMENTS

Foreword and Acknowledgements: Third Meeting of the International Neurotoxicology Association, ANTONIO MUTTI, LUCIO G. COSTA, LUIGI MANZO AND JOAN M. CRANMER .....	ix
---------------------------------------------------------------------------------------------------------------------------------------------------------------------	----

## II. SCREENING FOR NEUROTOXICITY IN HUMANS

Comparison of Screening Approaches. HUGH A. TILSON AND VIRGINIA C. MOSER .....	1
Cross Species Extrapolation in Neurotoxicology: Neurophysiological and Neurobehavioral Aspects. GERHARD WINNEKE .....	15
Neurotoxicology of PCBs and Related Compounds. WALTER J. ROGAN AND BETH C. GLADEN .....	27

## III. DEVELOPMENTAL NEUROTOXICITY

Models for the <i>In Vitro</i> Assessment of Neurotoxicity in the Nervous System in Relation to Xenobiotic and Neurotrophic Factor-Mediated Events. C. K. ATTERWILL, H. JOHNSTON AND S. M. THOMAS .....	39
Functional Neuroteratology of Drugs Acting on Adrenergic Receptors. MATTHIJS G. P. FEENSTRA .....	55
Animal Models for Human Behavioral Deficiencies During Development. JÜRGE ELSNER .....	65
Regulatory Developmental Neurotoxicity and Human Risk Assessment. ELAINE Z. FRANCIS .....	77

## IV. NEUROTOXICITY AND AGING

Reactive Oxygen Species: Relation to Aging and Neurotoxic Damage. STEPHEN C. BONDY .....	87
Neuroplasticity, the Aging Brain, and Alzheimer's Disease. THOMAS J. WALSH AND KAREN D. OPELLO .....	101

## V. CELLULAR AND SUBCELLULAR MECHANISMS OF NEUROTOXICITY

- Enhanced Expression of Glial Fibrillary Acidic Protein and the Cupric Silver Degeneration Reaction Can Be Used as Sensitive and Early Indicators of Neurotoxicity. J. P. O'CALLAGHAN AND K. F. JENSEN ..... 113
- Perturbation of Axonal Elemental Composition and Water Content: Implication for Neurotoxic Mechanisms. RICHARD M. LOPACHIN, CAROLYN M. CASTIGLIA AND ALBERT J. SAUBERMANN. .... 123
- Ca<sup>2+</sup>-Dependent Processes As Mediators of Neurotoxicity. M. ANTHONY VERITY ..... 139

## VI. *IN VITRO* NEUROTOXICOLOGY

- 2,5-Hexanedione Is a Potent Gliatoxin in *In-Vitro* Cell Cultures of the Nervous System. F. BOEGNER, W. GRÜNING, G. STOLTENBURG-DIDINGER, P. MARX AND H. ALTENKIRCH ..... 151
- Nerve Growth Factor (NGF) Receptors on Mast Cells: Effects of the Cholinergic Neurotoxin Ethylcholine Mustard Aziridinium Ion (ECMA). H. B. JOHNSTON AND C. K. ATTERWILL ..... 155
- Specific Neurotoxic Effects of Different Organic Solvents on Dissociated Cultures of the Nervous System. G. STOLTENBURG-DIDINGER, F. BOEGNER, W. GRÜNING, M. WAGNER, P. MARX AND H. ALTENKIRCH. .... 161
- Irradiation Neurotoxicity Assessed in Organotypic Cultures of Rat Hippocampus. G. HAMDORF, A. SHAHAR, J. CERVOS-NAVARRO, A. SKOBERLA AND A. SCHEFFLER ..... 165
- Neuronal Cell Cultures as a Model for Assessing Neurotoxicity Induced by Encephalitic Viruses. A. SHAHAR, H. SCHUPPER, S. LUSTIG, R. LEVIN, A. FRIEDMANN AND P. FUCHS ..... 171
- Glia Toxicity in Dissociated Cell Cultures Induced by Cyclosporine. G. STOLTENBURG-DIDINGER AND F. BOEGNER ..... 179
- In Vitro* Screening Batteries for Neurotoxicants. BELLINA VERONESI ..... 185

## VII. NEUROTOXICITY TO SENSORY ORGANS

- The Effects of 2,5-Hexanedione on Rods and Cones of the Retina of Albino Rats. B. BÄCKSTRÖM AND V. P. COLLINS ..... 199
- Evoked Potentials Are Modified by Long Term Exposure to Trichloroethylene. LUCIE BLAIN, PIERRE LACHAPPELLE AND STEPHANE MOLOTCHEV ..... 203
- Auditory Brainstem Evoked Potentials (BAEPs) in Lead-Exposed Workers. G. L. DISCALZI, F. CAPELLARO, L. BOTTALO, D. FABBRO AND A. MOCELLINI ..... 207
- Olfactory Threshold Shift Following Controlled 7-Hour Exposure to Toluene and/or Xylene. DONNA MERGLER AND BRIGITTE BEAUVAIS ..... 211

## VIII. NEUROBEHAVIORAL TOXICITY IN ANIMALS

- Comparative Studies on the Neuro- and Reproductive Toxicity of Acrylamide and its Epoxide Metabolite Glycidamide in the Rat. LUCIO G. COSTA, HAI DENG, CESARINA GREGOTTI, LUIGI MANZO, ELAINE FAUSTMAN, EMMA BERGMARK AND CARL J. CALLEMAN. . . . . 219
- Interactive Effects of Toluene and Hexane on Behavior and Neurophysiologic Responses in Fischer-344 Rats. GORDON T. PRYOR AND CHARLES S. REBERT . . . . . 225
- Increase in Rat Pup Ultrasonic Isolation Calls Induced by Lindane. S. RIVERA, C. SANFELIU, M. GARCÍA, F. COMELLAS AND E. RODRIGUEZ-FARRÉ . . . . . 235

## IX. NEUROTOXICITY IN HUMANS

- Neurobehavioral and Neuroendocrine Effects of Occupational Exposure to Perchloroethylene. C. FERRONI, L. SELIS, A. MUTTI, D. FOLLI, E. BERGAMASCHI AND I. FRANCHINI . . . . . 243
- Joint Effect of Gestational Age and Maternal Lead Exposure on Psychomotor Development of the Child at Six Years. GUY HUEL, PASCALE TUBERT, NADINE FRERY, THIERRY MOREAU AND JEAN DREYFUS . . . . . 249
- Long-Lasting Impairment of Neuroendocrine Response to Psychological Stress in Heroin Addicts. A. MUTTI, D. FOLLI, M. TH. VAN DER VENNE, A. BERLIN, G. GERRA, R. CACCAVARI, P. P. VESCOVI AND I. FRANCHINI . . . . . 255
- Calcium-Entry Blockers-Induced Parkinsonism: Possible Role of Inherited Susceptibility. A. NEGROTTI, S. CALZETTI AND E. SASSO . . . . . 261
- Correlations Between Subjective Disturbances Due to Acute Exposure to Organic Solvents and Internal Dose. A. SEEBER, E. KIRSSWETTER AND M. BLASZKEWICZ . . . . . 265
- Preclinical Neurophysiological Signs of Parkinsonism in Occupational Manganese Exposure. ARNE WENNBERG, MAUD HAGMAN AND LOTTA JOHANSSON . . . . . 271

## X. KINETICS AND DYNAMICS OF NEUROTOXIC CHEMICALS

- Accumulation of Inorganic Mercury in Lower Motoneurons of Mice. BJÖRN ARVIDSON . . . . . 277
- Inhibition of Muscarinic Receptor- and G-Protein-Dependent Phosphoinositide Metabolism in Cerebrocortical Membranes from Neonatal Rats by Ethanol. STEFANO M. CANDURA, LUIGI MANZO AND LUCIO G. COSTA . . . . . 281
- Effects of Diisopropyl Fluorophosphate on Muscarinic-M1-Receptors and on Receptor-Mediated Responsiveness of the Phosphatidyl-Inositol System in the Cerebral Cortex of Rats. ANNITA PINTOR, STEFANO FORTUNA, IRENA NALEPA AND HANNA MICHALEK . . . . . 289

Effects of Malaoxon on Phosphatidylinositol Signaling in Convulsing and Non-Convulsing Non-Pregnant and Pregnant Female Rats and Their Offspring. KAI M. SAVOLAINEN AND MAIIJA-RITTA R. HIRVONEN .....	295
Effect of $\gamma$ -Hexachlorocyclohexane and Its Isomers on Proto-Oncogene c-fos Expression in Brain. MONTSERRAT VENDRELL, JOSEP MARIA TUSELL AND JOAN SERRATOSA .....	301
CONTRIBUTORS.....	309
SUBJECT INDEX.....	313

---

**FRONT COVER:** The logo of the 3rd Meeting of the International Neurotoxicology Association – held in Salsomaggiore Terme, Parma, Italy, July 1-5, 1991 – is an original drawing by Franco Bolzonella (Fidia Research Laboratories, Abano Terme, Padua, Italy) representing a normal neuron before (white) and after (dark) exposure to a toxic chemical affecting the integrity of its dendritic processes.

# Increase in Rat Pup Ultrasonic Isolation Calls Induced by Lindane

S. RIVERA, C. SANFELIU, M. GARCÍA,\* F. COMELLAS† AND E. RODRÍGUEZ-FARRÉ

Department of Pharmacology and Toxicology, CSIC, c/Jordi Girona 18-26, 08034 Barcelona; \*Electronic Engineering Department, Polytechnic University of Catalonia, c/Jordi Girona s/n; 08034 Barcelona; †Applied Mathematics and Telematics Department, Polytechnic University of Catalonia, c/Jordi Girona s/n; 08034 Barcelona, Spain

**Abstract:** S. RIVERA, C. SANFELIU, M. GARCÍA, F. COMELLAS AND E. RODRÍGUEZ-FARRÉ. Increase in Rat Pup Ultrasonic Isolation Calls Induced by Lindane. *Neurotoxicology* 13(1):235-240, 1992. The neurotoxic agent lindane was tested for its ability to alter the rate of ultrasonic isolation calls of suckling rats. Doses that did not produce any sign of convulsant activity significantly increased the number of calls and the cumulative time of calling in male pups. At days 10 - 13 of age after a single dose of 20 mg/kg lindane, animals showed more than twice the control call values. After daily dosing with 10 mg/kg during the first week of age call increases also appeared. It is suggested that lindane has an anxiogenic effect mediated through its action on the benzodiazepine-GABA<sub>A</sub> receptor-chloride channel complex. © 1992 Intox Press, Inc.

**Key Words:** Ultrasonic Isolation Calls, Distress Calls, Suckling Rats,  $\gamma$ -Hexachlorocyclohexane, Lindane, Anxiety Testing

## INTRODUCTION

Mammalian vocalizations are a behavioral measure of emotional expression. One category of vocalization is the isolation call, given by infants when separated from their littermates or parents (Newman, 1988). Young rodents emit isolation calls in the ultrasonic range (Sales and Pye, 1974) which are a potent stimulus for maternal retrieval and care (Smotherman *et al.*, 1974). Calls increase in number and intensity along the 1st week of life to peak during the 2nd week, and abruptly decrease after the eye opening of the rat pup around day 15. This distress response is highly sensitive to environmental stimuli and pharmacological manipulations (Oswalt and Meier, 1975; Winslow and Insel, 1990).

Benzodiazepines (BDZ) are highly effective in decreasing vocalizations, probably via their anxiolytic action (Gardner and Budhram, 1987). Indeed, anxiogenic agents such as pentylenetetrazole (PTZ) selectively increase distress call emission (Insel *et al.*, 1986). Therefore, the GABA<sub>A</sub> receptor complex may play a physiological role in this basic emotional behavior.

Lindane ( $\gamma$ -hexachlorocyclohexane) is an hyperstimulant neurotoxic compound that binds to the picrotoxinin site on the chloride channel of the GABA-A receptor (Fishman and Gianutos, 1988). It has been suggested that this agent induces an anxiogenic response (Hijzen and Slangen, 1989; Llorens *et al.*, 1990). To test whether lindane would mimic anxiogenic agents in their effects on pup distress call behavior the present study was carried out.

Please send requests for reprints to Dr. Coral Sanfeliu, Department of Pharmacology and Toxicology, CSIC, c/Jordi Girona 18-26, E-08034 Barcelona, Spain.

Presented at the Third Meeting of the International Neurotoxicology Association, July 1-5, 1991 in Salsomaggiore Terme (Parma), Italy.

## MATERIAL AND METHODS

### Animals

Wistar rats (IFFA Credo, France) were used. One day after birth (day 1 of age) pups were randomized among the several litters. Each dam and its litter were individually housed under standard conditions.

### Treatment

Animals were administered vehicle (olive oil) or lindane (99.5% pure, Merck) by gavage (0.03ml/10 g body wt). A Silastic-R silicone elastic tubing was used for animals up to 8 days of age and conventional cannulae for older ones. Non-convulsant doses that were reported to produce some effects on motor behavior and cerebral 2-deoxyglucose uptake were used (Rivera *et al.*, 1990; submitted).

**Single Dosing.** 30 male rat pups aged 10 - 13 days (17 - 25 g body wt) were randomly assigned to the following treatments: vehicle, 10 mg/kg or 20 mg/kg lindane (10 animals/group). Testing was performed 1 hr later. Baseline values obtained in a pretrial performed immediately before administration showed no statistical differences between groups. Animals that failed to emit ultrasounds

in this preliminary testing were discarded from the study and replaced.

**Repeated Dosing.** 12 litters were randomly distributed among 4 daily treatments: vehicle or 10 mg/kg lindane, either during the 1st or the 2nd postnatal week. Only 2 males and 2 females from each litter were used in this study to avoid the litter effect (6 male and 6 female pups/group). Testing was performed on alternate days from the 2nd to the 20th day of age. Body weight was recorded twice a week.

### Apparatus and Procedure

Rat pups were individually tested. Litters were previously separated from their mothers and maintained in a cage with clean bedding warmed by infrared light. Each pup was gently put in a glass beaker inside a sound attenuated chamber. Recording of ultrasonic vocalizations started after 0.5 min of habituation time and lasted 2 min for the animals tested once and 3 min for those daily tested.

Vocalizations emitted were detected using a Knowles Electronic (BT1759-146FZ) microphone hanging 10 cm over the animal. Signals were fed through an amplifier (gain 120 dB) with a band-pass filter between 18 KHz (6th order) and 150 KHz (2nd order). A level detector digitalized the signals according as they were upper or lower than 2.2 V. This




FIG. 1. Changes in ultrasonic distress vocalizations after single administration of lindane. Bars represent the mean  $\pm$  SE of 10 rat pups of 10 - 13 days of age. A: Total number of calls during 2 min testing; B: Cumulative time of all calling emitted. (\* $p < 0.05$ ).






FIG. 2. Changes in ultrasonic distress vocalizations after daily administration of 10 mg/kg lindane during the 1st postnatal week. A: Total number of calls during 3 min testing. B: Cumulative time of calling. Groups are as follows: control males (—■—), lindane-dosed males (—●—), control females (---□---), and lindane-dosed female rats (---○---). Points represent the mean of 6 animals. For statistical evaluation, see Results.

voltage corresponded to a sound pressure level of 75 dB. Digital signals were sent to a PC computer through a port and recorded each msec during the whole testing time. Records were processed to obtain the number of calls and the cumulative duration of all calling emitted.

### Analysis of Data

One-factor ANOVA followed by Duncan's range test in the single dosing study and three-factor ANOVA (treatment, day and sex) in the repeated dosing experiments were performed.

## RESULTS

### Single Dose Experiment

Lindane 20 mg/kg increased ( $p < 0.05$ ) the number of emitted isolation calls and their cumulative time, as compared to the control group (Fig. 1).

### Repeated Dose Experiments

Animals dosed with lindane 10 mg/kg from day 1 to 7 of age showed a pattern of ultrasound emission different than the controls in the number of calls (ANOVA  $F_{(1,228)} = 6.16$ ,  $p =$

0.014) and in the total call time ( $F_{(1,228)} = 4.75$ ,  $p = 0.030$ ). The effect of sex was significant for the number of calls ( $F_{(1,228)} = 4.10$ ,  $p = 0.044$ ) but did not reach statistical significance for the total call time. If data were further analyzed after splitting them according to sex, ANOVA results for the treatment factor effect in males were  $F_{(1,109)} = 6.47$ ,  $p = 0.012$  and  $F_{(1,109)} = 4.14$ ,  $p = 0.044$ , respectively, for the number of calls and calling time. No significant differences were obtained for females (Fig. 2).

In the experiment where rat pups were dosed from day 8 to 14 of age, no statistical significance for treatment or sex effects were obtained (not shown).

The effect of the day of age at testing was highly significant for both parameters in both experiments ( $F_{(9,228)} = 9.38$ ,  $p < 0.001$  and  $F_{(9,228)} = 6.42$ ,  $p < 0.001$ , for the number of calls and total call time, respectively, in the 1st week administered animals;  $F_{(8,205)} = 5.98$ ,  $p < 0.001$  and  $F_{(8,205)} = 5.43$ ,  $p < 0.001$ , for the same parameters in the 2nd week dosing experiment).

Evolution of body weight was similar in control and lindane-treated animals (not shown).

## DISCUSSION

The age-related profile of ultrasonic isolation calls obtained in control pups was in gen-

eral agreement with that described in ontogenic studies (Sales and Pye, 1974).

Taken together, the results obtained indicate an increase of ultrasonic distress calls induced by lindane in isolated rat pups. Both parameters measured changed jointly and therefore the mean call duration was not modified by lindane treatment.

After a single dose of 20 mg/kg lindane calls were increased but 10 mg/kg had no effect. Insel *et al.* (1986) have found similar results with pentylenetetrazole (PTZ). Both compounds have been suggested to share common anxiogenic effects in adult rats in the plus-maze test (Llorens *et al.*, 1990). Hijzen and Slangen (1989) have also postulated an anxiogenic action of lindane in adult rats by startle response testing, after obtaining similar results with an inverse benzodiazepine (BDZ) agonist and opposite ones with the anxiolytic midazolam. Therefore, lindane-induced alterations of distress calling may be caused through a BDZ-modulated mechanism. This would be in agreement with the proposed mechanism of lindane action through the GABA system (Fishman and Gianutsos, 1988; Suñol *et al.*, 1989). Indeed, lindane antagonizes the GABA<sub>A</sub> receptor in the same way as PTZ does. Thus, the present results may provide further evidence for the role of the BDZ-GABA receptor-chloride channel complex in rodent attachment behavior.

Alterations of the age-related pattern of vocalizations were present at repeated doses of 10 mg/kg lindane when administered during the 1st week of age, but not during the 2nd one. Then, it may be postulated a higher sensitivity of the BDZ-binding site to a lindane-mediated effect during this early period. During the 1st postnatal week, for instance, the concentration of BDZ receptors increase from 20% to 60% of adult levels (Mallorga *et al.*, 1980). Distress isolation calls are the expression of an emotional behavior submitted to a multiple modulation which may include sex-related factors as observed in this study.

In conclusion, the present work supports an anxiogenic action of the neurotoxic agent lindane as suggested by previous experimental and clinical reports (Llorens *et al.*, 1990; Hijzen and Slangen, 1989; Kbare *et al.*, 1977).

## ACKNOWLEDGEMENTS

The authors wish to express gratitude to Ms. Carme Cleries for assistance with animal manipulations. This work was supported by grants 83/1648 from FIS and PB0202 from DGICYT.

## REFERENCES

- Fishman BE, Gianutsos G. CNS biochemical and pharmacological effects of the isomers of hexachlorocyclohexane (lindane) in the mouse. *Toxicol Appl Pharmacol* 1988; 93:146-153
- Gardner CR, Budhram P. Effects of agents which interact with central benzodiazepine binding sites on stress-induced ultrasounds in rat pups. *Eur J Pharmacol* 1987; 134:275-283
- Hijzen TH, Slangen JL. Effects of midazolam, DMCM and lindane on potentiated startle in the rat. *Psychopharmacology* 1989; 99:362-365
- Insel TR, Hill JL, Mayor RB. Rat pup ultrasonic isolation calls: Possible mediation by the benzodiazepine receptor complex. *Pharmacol Biochem Behav* 1986; 24:1263-1267
- Kbare SB, Rizvi AG, Shukta OP, Singh RRP, Perkash O, Misra VD, Gupta JP, Sethi PK. Epidemic outbreak of neuro-ocular manifestations due to chronic BHC poisoning. *J Assoc Physicians Ind* 1977; 25:215-222
- Llorens J, Tusell JM, Suñol C, Rodríguez-Farré E. On the effects of lindane on the plus-maze model of anxiety. *Neurotoxicol Teratol* 1990; 12:643-647
- Mallorga P, Hamburg M, Tallman JF, Gallagher DW. Ontogenetic changes in GABA modulation of brain benzodiazepine binding. *Neuropharmacology* 1980; 19:405-408
- Newman JD. Investigating the physiological control of mammalian vocalizations. In: *The Physiological Control of Mammalian Vocalization*, Newman JD, ed., New York, Plenum Press, 1988, pp 1-5
- Oswalt GL, Meier GW. Olfactory, thermal, and tactual influences on infantile ultrasonic vocalization in rats. *Develop Psychobiol* 1975; 8:129-135

- Rivera S, Sanfeliu C, Rodríguez-Farré E. Behavioral changes induced in developing rats by an early postnatal exposure to lindane. *Neurotoxicol Teratol* 1990; 12:591-595
- Sales G, Pye D. *Ultrasonic Communication by Animals*. London, Chapman and Hall, 1974, pp 149-201
- Smotherman WP, Bell RW, Starzec J, Elias J. Maternal responses to infant vocalizations and olfactory cues in rats and mice. *Behav Biol* 1974; 12:55-66
- Suñol C, Tusell JM, Gelpí E, Rodríguez-Farré E. GABAergic modulation of lindane ( $\gamma$ -hexachlorocyclohexane)-induced seizures. *Toxicol Appl Pharmacol* 1989; 100:1-8
- Wislow JT, Insel TR. Serotonergic and catecholaminergic reuptake inhibitors have opposite effects on the ultrasonic isolation calls of rat pups. *Neuropsychopharmacology* 1990; 3:51-59