

# Dynamics of the parabolic restricted three-body problem

Esther Barrabés<sup>a</sup>, Josep M. Cors<sup>b,\*</sup>, Mercè Ollé<sup>c</sup>

<sup>a</sup>*Universitat de Girona, Escola Politècnica Superior, 17071 Girona, Spain*

<sup>b</sup>*Universitat Politècnica de Catalunya, Escola Politècnica Superior de Manresa, Av. Bases de Manresa 61-73, 08242 Manresa, Spain. +34 938777247*

<sup>c</sup>*Universitat Politècnica de Catalunya, Escola Tècnica Superior d'Enginyeria Industrial de Barcelona, Av Diagona 647, 08028 Barcelona, Spain*

---

## Abstract

The main purpose of the paper is the study of the motion of a massless body attracted, under the Newton's law of gravitation, by two equal masses moving in parabolic orbits all over in the same plane, the planar parabolic restricted three-body problem. We consider the system relative to a rotating and pulsating frame where the equal masses (primaries) remain at rest. The system is gradient-like and has exactly ten hyperbolic equilibrium points lying on the boundary invariant manifolds corresponding to escape of the primaries in past and future time. The global flow of the system is described in terms of the final evolution (forwards and backwards in time) of the solutions. The invariant manifolds of the equilibrium points play a key role in the dynamics. We study the connections, restricted to the invariant boundaries, between the invariant manifolds associated to the equilibrium points. Finally we study numerically the connections in the whole phase space, paying special attention to capture and escape orbits.

*Keywords:* parabolic restricted three-body problem, invariant manifolds, final evolutions, global dynamics

*2010 MSC:* 70F07, 70F15

---

\*Corresponding author

*Email addresses:* barrabes@imae.udg.edu (Esther Barrabés), cors@eupm.upc.edu (Josep M. Cors), Merce.Olle@upc.edu (Mercè Ollé)

## 1. Introduction

Astronomy textbooks typically present galaxies as calm, solitary and majestic island worlds of glittering stars. However, the Hubble images support the well-known idea that galaxies are dynamic and energetic. In particular, the bridges and tails seen in some multiple galaxies are just relics of close encounters. The consequences of the brief but violent tidal forces have been studied by Toomre and Toomre in [16] and by Namboodiri et al. in [10] considering a simple-minded fashion: each encounter is considered to involve only two galaxies describing a roughly parabolic path.

This approach of the dynamics of the close encounters for two galaxies has been used, for example, by Condon et al. in [2], in the case of the galaxies UGC 12914 and 12915, or by Günthardt et al. in [7], for the system AM1003-435. The parabolic model has been also used in the study of the formation of planetary systems. Fagnier and Nelson, in [6], examine the effect of parabolic stellar encounters on the evolution of a Jovian-mass giant planet forming within a protoplanetary disc. Pfarzner et al., in [12], study the close encounter of two stars, one of them surrounded by a disc. More recently, in [14], Steinhausen et al. present a numerical investigation to study the effect of gravitational star-disc interactions on the disc-mass distribution, considering coplanar, prograde encounters on parabolic orbits.

A close approach of two galaxies (or stars surrounded by a disc) cause significant modification of the mass distribution or disc structure. Focussing just on one particle that initially stays in one galaxy (or around one star), after the close encounter, it can jump to the other galaxy or escape. One aim is to study the regions in the phase space where the particle remains or not around each galaxy. To perform this study, we consider a very simple model, the so called planar parabolic restricted three-body problem, which describes the motion of a massless particle submitted to the gravitational attraction of two masses -called primaries- that move in parabolic orbits around their common center of mass, when the primaries and the particle move in the same plane.

As far as the authors know, very little literature has been devoted to restricted three-body problems where the gravitational attraction of the primaries is non-periodic. That is, when the energy of the primaries is non-negative, so that they move in parabolic or hyperbolic orbits. Meyer and Wang, in [9],  
35 studied restricted isosceles problems (the three bodies are at the vertices of an isosceles triangle) when the energy of the primaries is non-negative and the infinitesimal mass moves in a line perpendicular to the orbital plane passing through the center of mass of the primaries. After that, Cors and Llibre, in [3] (see also the references therein), obtained a classification of the orbits in  
40 the parabolic case in terms of the asymptotic velocity of the infinitesimal mass when  $t \rightarrow \pm\infty$  and the number of times that the infinitesimal mass intersects the plane which contains the motion of the primaries. More recently, Alvarez et al., in [1], describe some features of the planar parabolic restricted three-body problem and show the existence of special types of motion. Finally, Faintich, in  
45 [5], considers the planar hyperbolic restricted three-body problem and applies the model to a hypothetical star-Sun-comet system to determine the effect of a stellar encounter on the orbit of the comet.

Being the previous works our original motivation, the main purpose of the paper is to continue and complete the results obtained in [1] and to describe  
50 the flow of the planar parabolic restricted problem (or simply *parabolic problem* along the paper). We consider in this work the simplest case, when the two primaries have equal masses. In particular, we will focus on the role of the invariant manifolds to explain how a particle can be captured around one primary or escape.

The paper is organized as follows: in Section 2, the equations of the motion  
55 of the parabolic problem are given both in an inertial system of coordinates and in a rotating and pulsating (synodic) frame. After that, in order to extend the flow of the system when the primaries are at infinity, the phase space is compactified in the time direction with the introduction of a suitable variable  
60  $\theta \in [-\pi/2, \pi/2]$ , and the so called *global* system is obtained. The flow when  $\theta = \pm\pi/2$  is invariant, obtaining the *upper* ( $\theta = \pi/2$ ) and the *lower* ( $\theta = -\pi/2$ )

boundary problems. In particular the equilibrium points and the homothetic solutions associated to them are considered and the gradient-like property of the equations is remarked. Finally the possible regions of motion (the so called Hill's regions) are described as well. Section 3 is divided in two parts. First, we analyze the final evolutions for the motion of the particle. We focus on *escape* and *capture* orbits in the inertial system of coordinates and we introduce a criterium (the so called *C-criterium*) that allows us to classify an orbit in the synodic system of coordinates. Second, we analyze the dynamics on the boundary problems. We will see that it is crucial to know what we call *map of heteroclinic connections* between invariant objects of the boundary system, that is, trajectories that start and end at a primary and/or an equilibrium point and/or the infinity. We recall some connections already known in [1] and we show the existence of new heteroclinic connections. Section 4 is devoted to numerical explorations. We present two strategies to find initial conditions of collision orbits, both forwards and backwards in time, called *connecting* orbits. Applying one or another strategy we obtain non symmetrical and symmetrical connecting orbits respectively. We will show that the equilibrium points and the invariant manifolds associated with them play a key role on the specific path of such connecting orbits and we will find orbits with close paths to triangular and/or collinear configuration during their trajectory.

## 2. Description of the problem and main features

In order to have a self contained paper, we present the equations of the motion of the problem and other main features. The details can be found in [1].

### 2.1. Equations of motion

Let us consider three bodies in an inertial (sidereal) reference system. Two of the bodies, called primaries, with masses  $m_1$  and  $m_2$ , move in parabolic orbits of the two body problem around their common center of mass. The third body is a particle of infinitesimal mass  $m_0$  that moves under the gravitational attraction

90 of the primaries without affecting them in the same plane of the motion of the primaries. The problem of the description of the motion of the particle is the planar parabolic restricted three body problem (simply *parabolic problem* along the paper). We will consider in this paper the case of two equal masses for the primaries, that in a suitable units means that we can take  $m_1 = m_2 = 1/2$ .

It is well known (see for example [4]) that, when the primaries move in a parabolic motion, the relative position vector from  $m_1$  to  $m_2$  is  $\mathbf{R} = (\sigma^2 - 1, 2\sigma)$  where  $\sigma = \tan(f/2)$ ,  $f$  is the true anomaly, their mutual distance is  $r = \sigma^2 + 1$ , and  $\frac{1}{\sqrt{2}}(t - T) = \sigma + \frac{\sigma^3}{3}$ , being  $T$  the time of passage at the pericenter (see Figure 1). Then, the equation of the motion of the particle in the inertial system of coordinates  $\mathbf{Z} = (X, Y)$  is

$$\ddot{\mathbf{Z}} = -\frac{1}{2} \frac{\mathbf{Z} - \mathbf{Z}_1}{|\mathbf{Z} - \mathbf{Z}_1|^3} - \frac{1}{2} \frac{\mathbf{Z} - \mathbf{Z}_2}{|\mathbf{Z} - \mathbf{Z}_2|^3}, \quad (1)$$

95 where  $\dot{\phantom{z}} = \frac{d}{dt}$  is the derivative with respect to the time  $t$ ,  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are the position vectors of the primaries, and we have assumed that the constant of gravitation  $G = 1$ .




Figure 1: Parabolic problem in an inertial (sidereal) system of reference.

After placing the common center of mass of the primaries at the origin, we perform two changes of variables. First, a standard rotating and pulsating (synodic) coordinate system  $\mathbf{z} = (x, y)$  is introduced, so that the primaries remain fixed along the  $x$ -axis at  $\mathbf{z}_1 = (-\frac{1}{2}, 0)$  and  $\mathbf{z}_2 = (\frac{1}{2}, 0)$  respectively. This change is done via the complex product

$$\mathbf{Z} = \mathbf{R} \cdot \mathbf{z}. \quad (2)$$

Second, we introduce the reparametrization of time  $\frac{dt}{ds} = \sqrt{2}r^{3/2}$ . After some straightforward computations, the variable  $\sigma$ , that gives the relative position of the primaries, can be expressed in terms of the new independent variable  $s$  as  $\sigma = \sinh(s)$ , and the equations of motion for the particle in the new coordinate system are

$$\begin{cases} \mathbf{z}' &= \mathbf{w}, \\ \mathbf{w}' &= A(s)\mathbf{w} + \nabla\Omega(\mathbf{z}) \end{cases} \quad (3)$$

where  $' = \frac{d}{ds}$  denote the derivative with respect to  $s$ ,

$$A(s) = \begin{pmatrix} -\tanh(s) & -4\operatorname{sech}(s) \\ 4\operatorname{sech}(s) & -\tanh(s) \end{pmatrix},$$

and  $\Omega$ , the potential function, is given by

$$\Omega(\mathbf{z}) = x^2 + y^2 + \frac{1}{\sqrt{(x - \frac{1}{2})^2 + y^2}} + \frac{1}{\sqrt{(x + \frac{1}{2})^2 + y^2}}.$$

Clearly, when the primaries tend to infinity along their parabolic orbits,  $t \rightarrow \pm\infty$  and the new time  $s$  also tends to  $\pm\infty$ . In order to extend the flow of the system when the primaries are at infinity, a new variable  $\theta$  is introduced through the change  $\sin(\theta) = \tanh(s)$ . With the new variables  $(\theta, \mathbf{z}, \mathbf{w})$  the system (3) becomes the following autonomous system

$$\begin{cases} \theta' &= \cos \theta, \\ \mathbf{z}' &= \mathbf{w}, \\ \mathbf{w}' &= -A(\theta)\mathbf{w} + \nabla\Omega(\mathbf{z}) \end{cases} \quad (4)$$

where

$$A(\theta) = \begin{pmatrix} \sin \theta & 4 \cos \theta \\ -4 \cos \theta & \sin \theta \end{pmatrix}.$$

Notice that the the original system (3) is defined for  $s \in (-\infty, \infty)$ , which corresponds to  $\theta \in (-\pi/2, \pi/2)$ , whereas the extended system (4) is defined also  
100 for  $\theta = \pm\pi/2$ , and it is invariant at the boundaries  $\theta = \pm\pi/2$ . Therefore the extended phase space is  $D = [-\pi/2, \pi/2] \times (\mathbf{R}^2 - \{(-1/2, 0), (1/2, 0)\}) \times \mathbf{R}^2$ .

From now on, we will call system (4) the *global* system, and we denote as *configuration space* the projection of  $D$  on to the  $(x, y)$  plane. The two invariant

systems will be denoted as the *upper* ( $\theta = \pi/2$ ) and *lower* ( $\theta = -\pi/2$ ) boundary problem respectively and the corresponding equations are

$$\begin{cases} \mathbf{z}' = \mathbf{w}, \\ \mathbf{w}' = \mp \mathbf{w} + \nabla \Omega(\mathbf{z}). \end{cases} \quad (5)$$

Finally, notice that the two changes

$$(s, \pm \frac{\pi}{2}, x, y, x', y') \rightarrow (-s, \mp \frac{\pi}{2}, x, y, -x', -y') \quad (6)$$

transform the equations of the upper/lower boundary problem to the equations of the lower/upper one. Thus, it is enough to study the flow and the dynamics of one of the boundary problems. We will describe properties and the dynamics  
105 of the upper boundary problem unless otherwise noted.

## 2.2. Symmetries

The global system (4) has the following two symmetries

$$(s, \theta, x, y, x', y') \rightarrow (-s, -\theta, x, -y, -x', y'), \quad (7)$$

$$(s, \theta, x, y, x', y') \rightarrow (s, \theta, -x, -y, -x', -y'). \quad (8)$$

The symmetry (7) implies that given a solution of the global system there exists another one which is symmetric with respect to  $y = 0$  in the  $(x, y)$  plane  
110 (reversing time and  $\theta$ ). The symmetry (8) implies that given a solution of the global system there exists another one which is symmetric with respect to the origin in the  $(x, y)$  plane.

The symmetries on the upper and lower boundary problems (5) are

$$(s, x, y, x', y') \rightarrow (s, x, -y, x', -y'), \quad (9)$$

$$(s, x, y, x', y') \rightarrow (s, -x, -y, -x', -y'). \quad (10)$$

For each solution of a given boundary problem there exists another one of the  
115 same problem that is symmetric with respect to  $y = 0$  in the  $(x, y)$  plane (using (9)) and another one symmetric with respect to the origin (using (10)).

We emphasize that the symmetries (8) and (10) are specific for the parabolic problem with equal masses, whereas (7) and (9) apply for the parabolic problem for any value of the mass parameter.

120 *2.3. Equilibrium points*

The equilibrium points of the global system (4) are given by

$$\theta = \pm\pi/2, \quad \mathbf{w} = 0, \quad \nabla\Omega(\mathbf{z}) = 0.$$

Therefore, all the equilibrium points of the global system are at the upper and lower boundary problems (5). Moreover, the potential function  $\Omega$  satisfies

$$\Omega(\mathbf{z}) = 2\bar{\Omega}(\mathbf{z}), \quad (11)$$

where  $\bar{\Omega}(\mathbf{z})$  is the potential function of the circular restricted three-body problem in rotating coordinates (see [15]). Thus, the equilibrium points of the parabolic problem in each boundary coincide with the classical equilibrium points of the restricted circular three-body problem: three collinear and two triangular. We denote by  $L_i^+$  and  $L_i^-$ ,  $i = 1, \dots, 5$ , the equilibrium points for  $\theta = \pi/2$ , and  $\theta = -\pi/2$  respectively. Note that due to the symmetries (9) and (10)  $L_1^\pm$  and  $L_5^\pm$  are opposite to  $L_3^\pm$  and  $L_4^\pm$  respectively. See Table 1.

Linearizing the upper boundary problem, the eigenvalues associated to the collinear equilibrium points  $L_i^+$ ,  $i = 1, 2, 3$ , are

$$\lambda_k^1 > 0, \quad \lambda_k^2 < 0, \quad \lambda_k^{3,4} = -\frac{1}{2} \pm ib_k, \quad k = 1, 2, 3.$$

Thus, the collinear equilibrium points have an unstable manifold  $W^u(L_i^+)$  of dimension 1 and a stable manifold  $W^s(L_i^+)$  of dimension 3. Due to symmetry (6), in the lower boundary problem the dimensions are the contrary (the unstable is of dimension 3, the stable of dimension 1). In the case of the triangular equilibrium points  $L_i^+$ ,  $i = 4, 5$ , the linearization gives the eigenvalues

$$\lambda_k^{1,2} > 0, \quad \lambda_k^{3,4} < 0, \quad k = 4, 5.$$

Thus, both invariant manifolds associated to the triangular equilibrium points, the unstable and the stable one, are of dimension 2.

130 Considering the equilibrium points in the global system (4), the dimension of  $W^u(L_i^-)$  and  $W^s(L_i^+)$ ,  $i = 1, \dots, 5$  increases in one. See Table 1.

	$L_{1,3}^+$	$L_2^+$	$L_{4,5}^+$
$(x_i, y_i)$	$(\mp 1.198406145, 0)$	$(0, 0)$	$(0, \pm\sqrt{3}/2)$
$C(L_i^+) = C_i$	6.91359245	8	5.5
$\dim(W^u)$	1		2
$\dim(W^s)$	4		3

Table 1: Equilibrium points of the upper boundary problem: position, value of the Jacobi function (see Section 2.5), and dimension of the invariant manifolds in the global system.

#### 2.4. Homothetic solutions

Besides the equilibrium points, the simplest solutions of the global system (4) are the five homothetic solutions connecting the equilibrium points  $L_i^-$  with  $L_i^+$ , and belonging to  $W^u(L_i^-) \cap W^s(L_i^+)$ :

$$\theta(s) = \arcsin(\tanh(s)), \quad \mathbf{z}(s) = (x_i, y_i), \quad \mathbf{w}(s) = 0, \quad i = 1, \dots, 5$$

where the coordinates  $(x_i, y_i)$  are given in Table 1.

Clearly these five homothetic solutions in the rotating-pulsating coordinate system are homographic solutions of the original coordinate system (1). They are solutions in which the three bodies keep the same configuration all the time: either the three bodies lie in a line (collinear configuration) or they lie at the vertices of an equilateral triangle (triangular configuration).

#### 2.5. Jacobi function and Hill's regions of motion

We consider the analogous function to the Jacobi constant of the circular restricted three-body problem (see [15]), that we call, by similarity, the Jacobi function:

$$C = 2\Omega(\mathbf{z}) - |\mathbf{w}|^2. \quad (12)$$

In the parabolic problem, however,  $C$  is not constant along the solutions of the global system (4), because (see Proposition 1 in [1])

$$\frac{dC}{ds} = 2 \sin \theta |\mathbf{w}|^2. \quad (13)$$

140 Notice that  $C$  does not depend on  $\theta$  explicitly, while its derivative does. Furthermore, the Jacobi function has a piecewise monotone behavior along the solutions of the global system (except at the homothetic ones, where the Jacobi function is constant). This behavior is known as gradient-like property. More precisely, along any solution of the global system, when  $\theta \in [-\pi/2, 0]$  ( $s \leq 0$ ) the function  
145  $C$  decreases, whereas for  $\theta \in [0, \pi/2]$  ( $s \geq 0$ ) the function  $C$  increases. In the boundary problems, the function  $C$  is monotone (for  $\theta = -\pi/2$ ,  $C$  decreases and for  $\theta = \pi/2$ ,  $C$  increases). A particular and important consequence of this gradient-like property is that there cannot exist periodic orbits.

From (12) we obtain the so called Hill's regions, that is, the allowed regions of motion in the configuration space. For a fixed value of  $C$ , we consider the zero velocity set,  $V_0(C)$ , in the configuration space defined by

$$V_0(C) = \{\mathbf{z} \mid 2\Omega(\mathbf{z}) = C\}.$$

It can be shown that  $V_0(C)$  is a set of closed curves (called zero velocity curves or  
150 zvc). Due to (11), the topology of the zvc is the same as in the circular restricted three body problem (see [15]). In order to describe them, let  $C_i = C(L_i^\pm)$ ,  $i = 1, \dots, 5$  be the value of the Jacobi function at the equilibrium points. We observe that  $C_1 = C_3$ ,  $C_4 = C_5$ , and in Table 1 the (approximate) values of  $C_2$ ,  $C_3$  and  $C_4$  are given. We plot in Figure 2 the zvc and the forbidden regions of  
155 motion (shaded regions) for different fixed values of  $C$ .

As the Jacobi function varies with time, the Hill's regions evolve also with time. We describe the evolution of the geometry of the zvc as the value of the Jacobi function increases. For  $C < C_4 = C_5$ , the set  $V_0(C)$  is empty and the motion is possible everywhere. For  $C_4 \leq C < C_{1,3}$ , the zvc are two closed  
160 curves surrounding the triangular equilibrium points (Figure 2, a) and b)), which increase in size as  $C$  increases. For  $C_{1,3} < C < C_2$ , the zvc still have two components, one around the primaries and  $L_2^\pm$  and an exterior one enclosing all the equilibrium points (Figure 2, c)), so there are two regions where the motion is admissible. Finally, for  $C > C_2$ , the interior component of the zvc becomes  
165 two unconnected curves, each one surrounding one primary (see Figure 2, d)).




Figure 2: Zero velocity curves and the forbidden (shaded) regions of motion in configuration space for fixed values of the Jacobi function. From left to right:  $C = 6$ ,  $C = 6.9 < C_1 = C_3$ ,  $C_1 < C < 7$  and  $C_2 < C < 8.1$ .

Taking into account the gradient-like property (13), the piecewise monotone behavior of the Jacobi function for  $s \leq 0$  and  $s \geq 0$ , and the geometry of the zvc, we have that following a solution forwards in time, the regions of admissible motion grow when  $\theta \in [-\pi/2, 0]$  and shrink when  $\theta \in [0, \pi/2]$ .

### 170 3. Dynamics of the parabolic problem

In order to describe the dynamics of the parabolic problem, we will focus on two aspects: the final evolutions when time tends to infinity, and the existence of heteroclinic connections between equilibrium points.

We will see that the behavior of the trajectories as time tends to infinity is  
 175 rather simple, as it cannot be otherwise in a problem where periodic orbits do not exist. There are only three possible final evolutions for a trajectory: or it escapes far away from the primaries, or it tends to one of them, or it tends to one of the equilibrium points. And by the symmetry of the problem, the origins of the trajectories are the same. Despite this simplicity in the final evolution  
 180 of the orbits, the problem exhibits a richness in the intermediate stages due to the existence of invariant manifolds associated to the homographic solutions. The invariant manifolds of codimension 1 are the natural frontiers in the phase space,  $Int(D)$ , that separate the different types of orbits. Furthermore, the heteroclinic connections between the equilibrium points allow the existence of  
 185 orbits with passages close to collinear and/or equilateral configurations. Thus,

in order to fully understand the global system flow, all possible heteroclinic connections concerning the equilibrium points must be determined.

First, in order to have a tool to classify the orbits depending on their final evolution, we give a criterium that allows to decide when an orbit is captured  
 190 by one of the primaries or when it tends to infinity. Next, we show the existence of some heteroclinic connections and complete the diagram of heteroclinic connections given in [1].

### 3.1. Final evolutions

We focus on the ultimate behavior of the motion of a particle when the  
 195 time tends forwards to infinity, i.e. the  $\omega$ -limit (see, for example, [11]) of the solutions of the parabolic problem. Due to the symmetries of the problem, the global dynamics of the problem backwards in time (and the  $\alpha$ -limit of the solutions) can be obtained from the analysis of the dynamics forwards in time.

We focus on *escape* and *capture* orbits. Essentially, an escape orbit will be a  
 200 path along which the particle moves away from both primaries and their center of mass, whereas a capture orbit will be a trajectory along which the particle approaches one of the primaries, in the sense that while the primary tends to infinity along its parabolic orbit, the particle follows the primary (spinning around it) at a bounded distance. From Painleve's theorem (see for example Chapter  
 205 4 in [13]), if the right maximal interval of existence  $[0, \beta)$  is finite, a collision in finite time occurs. That case can be considered as capture. For this reason, from now on we will only consider orbits such that its right maximal interval of existence is  $[0, \infty)$ . All the definitions and results are given considering that we can take the limit when time ( $t$  and  $s$ ) tends to infinity, so that  $\theta$  tends to  $\pi/2$ .

210 The definitions of capture and escape that we use in the present paper are the following ones.

**Definition 1.** *Let  $\mathbf{Z}(t)$  be a solution of the parabolic problem given by equations (1). We say that*

- *it is a capture orbit around the primary of mass  $m_i$ , for  $i = 1$  or  $2$ , if*  
 215  $\limsup_{t \rightarrow \infty} |\mathbf{Z}(t) - \mathbf{Z}_i(t)| \leq K$ , *for some constant  $K$ ;*

- it is an escape orbit if  $\limsup_{t \rightarrow \infty} |\mathbf{Z}(t)| = \infty$  and  $\limsup_{t \rightarrow \infty} |\mathbf{Z}(t) - \mathbf{Z}_i(t)| = \infty$  for  $i = 1$  and  $2$ .

It is important to point out that the definition of capture and escape is done in the inertial reference system. From the change of variables (2) we have that

$$|\mathbf{Z} - \mathbf{Z}_i| = r|\mathbf{z} - \mathbf{z}_i|,$$

where the distance between the primaries  $r \nearrow \infty$  as time tends to infinity. Clearly, to be a capture orbit it is necessary that  $\lim_{s \rightarrow \infty} |\mathbf{z}(s) - \mathbf{z}_i| = 0$ , but that condition is not sufficient. On the contrary, to be an escape orbit it is enough that the  $\liminf_{s \rightarrow \infty} |\mathbf{z}(s)| \geq K$ , for  $K$  big enough to ensure that the particle does not approach the primaries when  $s \nearrow \infty$ . The aim is to have a criterium that allows to check whether a solution satisfies these conditions in the synodic reference system.

We introduce the concept of a collision orbit in the synodic reference system.

**Definition 2.** Let  $\gamma(s) = (\theta(s), \mathbf{z}(s), \mathbf{w}(s))$ ,  $s \in [0, \infty)$ , be a solution of the global system (4). We say that it is a collision orbit if  $\liminf_{s \rightarrow \infty} |\mathbf{z}(s) - \mathbf{z}_i| = 0$ , for some  $i = 1, 2$ .

Next, our aim is to prove that in the synodic system, the solutions of the global system can be classified in three classes: collision orbits, trajectories that go to infinity or trajectories tending to an equilibrium point. The first ones are candidates to be capture orbits, while the second ones are escape orbits.

**Lemma 1.** Let  $\gamma(s) = (\theta(s), \mathbf{z}(s), \mathbf{w}(s))$ ,  $s \in [0, \infty)$ , be a solution of the global system (4). Suppose that there exist positive constants  $M$  and  $\delta$  such that

$$\liminf_{s \rightarrow \infty} |\mathbf{z}(s)| \leq M, \quad \text{and} \quad \limsup_{s \rightarrow \infty} |\mathbf{z}(s) - \mathbf{z}_i| \geq \delta > 0,$$

$\forall i = 1, 2$ . Then, the  $\omega$ -limit of the solution is an equilibrium point.

*Proof.* From the three limits of the hypothesis, there exists a sequence  $\{s_k\}_{k \in \mathbb{N}}$ ,  $s_k \nearrow \infty$ , such that

$$|\mathbf{z}(s_k)| \leq M, \quad \text{and} \quad |\mathbf{z}(s_k) - \mathbf{z}_i| \geq \delta, \quad \forall i = 1, 2 \text{ and } k \in \mathbb{N}.$$

On one hand,  $\theta(s_k) \rightarrow \pi/2$ , so  $\{C(\gamma(s_k))\}_{k \in \mathbb{N}}$  is an increasing sequence. On the other hand,  $\{\Omega(\mathbf{z}(s_k))\}_{k \in \mathbb{N}}$  is bounded. Thus,  $\{|\mathbf{w}(s_k)|\}_{k \in \mathbb{N}}$  must be also bounded, and there exists the limit

$$\lim_{k \rightarrow \infty} C(\gamma(s_k)) = C_\infty < \infty.$$

In fact, as  $C(\gamma(s))$  increases for  $s > 0$ , the whole  $\lim_{s \rightarrow \infty} C(\gamma(s)) = C_\infty$ .

Furthermore,  $\{\gamma(s_k)\}_{k \in \mathbb{N}}$  is contained in a compact set, so there must exist a convergent subsequence. To simplify notation, we suppose that the sequence is convergent, so

$$\lim_{k \rightarrow \infty} \gamma(s_k) = \mathbf{q},$$

235 where  $\mathbf{q} = (\pi/2, \mathbf{z}_q, \mathbf{w}_q)$ , belongs to the  $\omega$ -limit of  $\gamma(s)$ , and  $|\mathbf{z}_q| \leq M$  and  $|\mathbf{z}_q - \mathbf{z}_i| \geq \delta, \forall i = 1, 2$ . Denote by  $\Gamma$  the solution of the global system through  $\mathbf{q}$ .

Let  $B$  a ball of center  $\mathbf{q}$  and radius  $\delta/2$ , so that the vector field of equations (4) does not have any singularity in  $B$ . Then, the set  $\Gamma \cap B$  is also contained  
240 in the  $\omega$ -limit of  $\gamma(s)$ . By continuity, the Jacobi function along  $\Gamma \cap B$  must be constant equal to  $C_\infty$ . And the only solutions of the global system with a constant value of the Jacobi constant are the equilibrium points.

Finally, the  $\omega$ -limit must be a connected set, so it can only contain one equilibrium point.  $\square$

245 **Corollary 1.** *Let  $\gamma(s)$  be a collision orbit. Then  $\lim_{s \rightarrow \infty} |\mathbf{z}(s) - \mathbf{z}_i| = 0$ , for some  $i = 1, 2$ .*

As a consequence, we obtain the following results.

**Proposition 1. (Final evolutions)** *Let  $\gamma(s) = (\theta(s), \mathbf{z}(s), \mathbf{w}(s))$ ,  $s \in [0, \infty)$ , be a solution of the global system (4). Then, either it is a collision orbit, or  
250  $\lim_{s \rightarrow \infty} |\mathbf{z}(s)| = \infty$  or its  $\omega$ -limit is an equilibrium point.*

Using the above statements, we can give the following criterium, named the  $C$ -criterium because the gradient-like property of the Jacobi function  $C$  is the main key.

**Proposition 2. (*C-criterion*)** Let  $\mathbf{q} \in \text{Int}(D)$  with  $\theta \geq 0$ , and  $\gamma(s) =$   
 255  $(\theta(s), \mathbf{z}(s), \mathbf{w}(s))$ ,  $s \in [0, \infty)$ , the solution of the global system (4) through  $\mathbf{q}$ .  
 Then,

- (i) if for some time  $s_0$  the value of the Jacobi function  $C(\gamma(s_0)) > C_2$  and  
 $\mathbf{z}(s)$  is located in one of the bounded components of the Hill's region, then  
 it is a collision orbit;
- 260 (ii) if for some time  $s_0$  the value of the Jacobi function  $C(\gamma(s_0)) > C_3$  and  
 $\mathbf{z}(s)$  is located in the unbounded component of the Hill's region, then it is  
 an escape orbit.

*Proof.* Suppose  $\gamma(s)$  satisfies the hypothesis of the first item. If it was not a  
 collision orbit, by Lemma 1 its  $\omega$  limit should be an equilibrium point, which is  
 265 not possible because in the bounded component of the Hill's region limited by  
 $V_0(C)$ , for  $C > C_2$  there are no equilibrium points.

In the second assumption, suppose that  $\liminf_{s \rightarrow \infty} |\mathbf{z}(s)| \leq K$  for some  
 constant  $K$ . Again, applying the Lemma 1 we will get a contradiction. Thus,  
 the limit

$$\lim_{s \rightarrow \infty} |\mathbf{z}(s)| = \infty$$

and then it is an escape orbit. □

In Theorem 4 of [1], the authors give a similar criterium (with the same  
 name) but with some important differences. The first one is that the conclusion  
 of the theorem is that the orbit is a collision orbit or its  $\omega$ -limit is an equilibrium  
 point, but they forgot to include the hypothesis that the particle must be in  
 the bounded component of the Hill's region. So, they do not consider the  
 escape orbits. Another important issue is that the authors use in the proof the  
 fact that the Jacobi function  $C(\gamma(s)) \nearrow \infty$  for orbits that do not tend to an  
 equilibrium point, although, as far as we know, this fact is not proved. Finally,  
 the authors use the result to prove the existence of capture orbits. But we want  
 to remark that whereas escape orbits can be detected studying their behavior  
 in the synodic reference system, it is not sufficient to know that an orbit is a

collision orbit to ensure that it is a capture orbit. In fact, in order to have a capture orbit it would be necessary to see if

$$\limsup_{t \rightarrow \infty} |\mathbf{Z}(t) - \mathbf{Z}_i(t)| = \limsup_{s \rightarrow \infty} \cosh^2(s) |\mathbf{z}(s) - \mathbf{z}_i| \leq K$$

for some constant  $K$ , which means that we would be able to check if  $|\mathbf{z}(s) - \mathbf{z}_i| = \mathcal{O}(e^{-2s})$ .

270 *3.2. Dynamics of the upper boundary problem*

Once we have classified the possible final evolutions of the solutions of the parabolic problem, we focus now on the dynamics of the upper (similarly lower) boundary problem. More precisely, we will study all possible connections involving the equilibrium points. These connections will provide different specific  
275 ways of tending to a final evolution. Taking also into account the homographic solutions, a rich variety of solutions of the global system with close passages to collinear and triangular configurations will be obtained.

In [1] some heteroclinic connections are studied. Our aim is to add some other connections to the diagram given by the authors. In Figure 3 we show  
280 the connections proved in ([1]) with dotted (black) lines, as well as the new ones by dashed (green) and continuous (red) lines. The equilibrium points on the diagram are located such that from bottom to top the value of the Jacobi function increases, and the Hill's regions shrink.

First we analyze which connections of the diagram are admissible. On one  
285 hand, recall that the Jacobi function increases when  $s$  increases, so it is not possible to connect  $L_i^+$  with  $L_j^+$  if  $C(L_i^+) \leq C(L_j^+)$ . On the other hand, the connections given in [1] were proved using the invariance of the planes  $y = y' = 0$  and  $x = x' = 0$ , and the study of the flow restricted to each plane. From that, it is clear that  $W^u(L_i^+)$ ,  $i = 1, 2, 3$ , which are of dimension 1, are contained in  
290 the  $y = 0$  axis. Then, neither  $L_1^+$ , nor  $L_3^+$ , connects with  $L_2^+$ . For the same reason, there is no connection from  $L_1^+$  to the second primary or from  $L_3^+$  to the first primary. And similarly, there is no connection from  $L_2^+$  to infinity.




Figure 3: Map of connections in the upper boundary problem. The dotted (black) lines are connections already known in [1]. The dashed (green) and continuous (red) lines are new connections.

Therefore, all possible connections starting at one of the collinear equilibrium points are already known and drawn in Figure 3. Looking at the possible connections emanating from  $L_4^+$  and  $L_5^+$ , we want to show that the following heteroclinics exist:

- (i) from  $L_{4,5}^+$  to  $\infty$  (dashed green lines),
- (ii) from  $L_{4,5}^+$  to  $L_{1,3}^+$  (continuous red lines),
- (iii) from  $L_{4,5}^+$  to the primaries (continuous red lines).

As we will see, we use essentially the behavior of the zero velocity sets  $V_0(C)$  and the invariant manifolds of the equilibrium points to show the above heteroclinic connections emanating from the triangular equilibrium points. A similar methodology can not be applied in the case of connections *arriving* at the triangular points. When exploring the behavior of the orbits of the  $W^s(L_{4,5}^+)$  backwards in time, the Jacobi function decreases to values lower than 4, and all the configuration space is available. So, the zvc do not play any role. For that reason, we have not explored the connections from one primary, or infinity, to the triangular equilibrium points.

We start justifying connection (i): from  $L_{4,5}^+$  to infinity (dashed green lines

310 in Figure 3). They can be proved immediately from the fact that the plane  
 ( $y, y'$ ) is invariant, and the equilibrium points  $L_{4,5}^+$  are saddle. That implies, in  
 particular, that there exist orbits of the invariant manifolds  $W^u(L_{4,5}^+)$  that lie  
 on the  $x = 0$  axis, one branch of them tending to  $L_2^+$ , the other one escaping to  
 infinity.

315 Next, we want to show numerically the existence of connection (ii): from  $L_4^+$   
 to  $L_3^+$ . By symmetry, all the other connections from  $L_{4,5}^+$  to  $L_{1,3}^+$  are obtained  
 (red continuous lines in Figure 3). We examine the behavior of the invariant  
 manifold  $W^u(L_4^+)$  forwards in time. In order to do so, we consider an approx-  
 imated parametrization of the invariant manifold, which is of dimension two,  
 320 and we propagate the initial conditions given by the parametrization forwards  
 in time until a certain section close to  $L_3^+$ . To obtain such intersections with  
 a good accuracy, the linear approximation of the parametrization is not good  
 enough. We have considered the fourth order approximation (see [8] for details  
 on the computation of the invariant manifold of an equilibrium point up to a  
 325 desired order).

Given a value of the Jacobi function  $C = C^*$ , we define the section

$$\Sigma_{C^*} = \{(\mathbf{z}, \mathbf{w}) \mid C(\mathbf{z}, \mathbf{w}) = C^*\},$$

and the intersection of the invariant manifold with that section is the set

$$W_{C^*}^u(L_i^+) = W^u(L_i^+) \cap \Sigma_{C^*}.$$

The invariant manifold  $W^u(L_4^+)$  is of dimension 2, so for values of  $C^*$  greater  
 but close to  $C_4$  (on the upper boundary problem, the Jacobi function increases),  
 the set  $W_{C^*}^u(L_4^+)$  is a simple closed curve (see Figure 4).

We want to study the evolution of the sets  $W_{C^*}^u(L_4^+)$  as  $C^*$  approaches  
 330  $C_3 \simeq 6.91359245$ . On one hand, the zero velocity set  $V_0(C^*)$  has two components  
 (one in  $y > 0$ , the other one in  $y < 0$ ) that approach to each other at the point  
 $L_3^+$  as  $C^* \nearrow C_3$  (see Figure 2 a) and b)). On the other hand, the projection  
 in configuration space of  $W_{C^*}^u(L_4^+)$  is a closed curve that surrounds one of the  
 components of  $V_0(C^*)$  (the one in the halfplane  $y > 0$ ), and, when  $C^* \nearrow C_3$ , this




Figure 4: Upper boundary problem: projection of  $W_{C^*}^u(L_4^+)$  in configuration space for some values of  $C^* \in (C_4, C_3)$ .

335 curve is trapped between the two components of  $V_0(C^*)$ . So at  $C^* = C_3$ , the two components of the zero velocity curve and the set  $W_{C_3}^u(L_4^+)$  must coincide at  $L_3^+$ , and at least one orbit on the invariant manifold must end up at the equilibrium point  $L_3^+$  (see Figure 5 and 6).




Figure 5: Upper boundary problem: projection of  $W_{C^*}^u(L_4^+)$  (red) in configuration space and  $Z_0(C^*)$  (black) for the values  $C^* = 6.9$  (left) and  $C^* = 6.913$  (right)

Finally, we observe that when we consider the points of the subset of  $W_{C_3}^u(L_4^+)$  340 belonging to the bounded component of the Hill's region, the associated trajectories will end up at a collision with one primary except the ones that tend to  $L_2^+$ . So the heteroclinic connections (iii) from  $L_{4,5}^+$  to the primaries follow.




Figure 6: Upper boundary problem: projection of  $W_{C^*}^u(L_4^+)$  (blue) in configuration space and  $Z_0(C^*)$  (black) for the values  $C^* = 6.913$  and  $C^* = 6.9136$ .

#### 4. Numerical results

As stated before, the simplest solutions in the parabolic problem are the  
 345 homographic solutions, which remain fixed in the  $(x, y)$  plane. Among all other  
 possible orbits or trajectories in the parabolic problem, we are interested, due to  
 astronomical reasons, in ones that go from primary to primary (either the same  
 one or not) that is, capture orbits forwards and backwards in time. As explained  
 in Section 3, we have a criterium to determine when a solution of the parabolic  
 350 problem in the synodic system satisfies the necessary condition to be a capture  
 orbit in the sidereal system. That is, when an orbit in the synodic system is of  
 collision forwards in time. We want to use that criterium to find sets of initial  
 conditions corresponding to collision orbits both forwards and backwards. We  
 call them *connecting* orbits. Our purpose is to present strategies that allow us  
 355 to find initial conditions of such type of motion. We will also discuss strategies  
 to find connecting orbits with a priori desired path with close approaches to  
 triangular or/and collinear configurations.

We will see that the two strategies presented are based on the knowledge  
 of the flow on the boundary manifolds,  $\theta = \pm\pi/2$ , and on the invariant mani-  
 360 folds associated to their equilibrium points, and whereas the first strategy gives  
 rise to (typically) non symmetrical connecting orbits, with the second one only  
 symmetrical connecting orbits are obtained.

4.1. *Connecting orbits with passages close to collinear or triangular configurations*

365 We consider from now on the global system given by (4) and we focus our attention on connecting orbits with a passage close to a collinear or a triangular configuration. We say that a connecting orbit for the global system is of type  $m_i - L_k - m_j$ , for  $i, j \in \{1, 2\}$  and  $k \in \{1, \dots, 5\}$ , if it is a collision orbit with  $m_i$  backwards in time, collision orbit with  $m_j$  forwards in time, and along its  
370 trajectory it has a close approach to the homothetic solution generated by  $L_k^\pm$  –in short, we will say that the orbit has a close passage to  $L_k$ , although strictly speaking there are no equilibrium points in  $Int(D)$ –. This implies that at some time the position of the particle and the primaries is close to a collinear or triangular configuration. We observe that, due to the connection diagram of each boundary problem ( $\theta = \pm\pi/2$ ) discussed in the previous section, it is  
375 natural to find connecting orbits that have several passages close to different  $L_m$ ,  $m \in \{1, \dots, 5\}$ .

The procedure to find initial conditions that lead to (typically non-symmetric) connecting orbits of type  $m_i - L_k - m_j$  is described as follows. Consider an invariant manifold of an equilibrium point of the lower boundary,  $W^{s,u}(L_k^-)$  for  
380 some  $k \in \{1, \dots, 5\}$ . We take initial conditions  $(\mathbf{z}_0, \mathbf{w}_0)$  on  $W^{s,u}(L_k^-)$  but with a value of  $\theta_0$  at a certain distance of  $-\pi/2$ ,  $\theta_0 = -\pi/2 + \delta$  with  $\delta > 0$ . We integrate this initial condition for the global system forwards and backwards in time for  $\theta \in [-\pi/2 + \epsilon_1, \pi/2 - \epsilon_2]$ , with  $0 < \epsilon_1, \epsilon_2$ , and  $\epsilon_1 < \delta$ . The key point is  
385 to find suitable initial conditions  $(\theta_0, \mathbf{z}_0, \mathbf{w}_0)$  such that we can guarantee that for the global flow, the associated orbit is a connecting orbit having a close path to  $L_k$  (or even to more equilibrium points). In general, we take initial conditions on  $W^s(L_k^-)$  and a small value of  $\delta$  (usually of order  $10^{-3}$ ). Starting close to the lower boundary problem and integrating backwards, we can use the  
390 knowledge of the behavior of the invariant manifolds in that boundary problem to obtain collision orbits with a desired primary. To select the appropriate initial conditions corresponding to collision orbits forwards in time, we apply the  $C$ -criterion. It is also possible to start with  $W^u(L_i^-)$  (that is the case of the

examples shown in Figure 8). In that case, we have considered bigger values  
 395 of  $\delta$  (of order 0.05), so that the initial conditions are not so close to the lower  
 boundary problem. In this case is not possible to use the knowledge of the  
 behavior of that invariant manifold on the lower boundary problem, and the  
 $C$ -criterium is also used backwards in time to select the initial conditions that  
 correspond to collision orbits.

400 Finally, the same strategy can be applied starting at initial conditions close  
 to the upper boundary manifold.

We show some particular examples of different connecting orbits visiting one  
 or more equilibrium points. The plots in configuration space  $\mathbf{z}$  show the location  
 of the equilibrium points with crosses (in fact the projection in configuration  
 405 space of the homothetic solutions). In the plots using the inertial coordinates  
 $\mathbf{Z}$ , we show the positions of the primaries (along their parabolic orbits) and the  
 particle at three different times  $t_1 < t_2 < t_3$ . The time  $t_2$  has been chosen such  
 that the trajectory of the particle is close to a collinear or triangular configura-  
 tion. The marks used to show the location of the bodies at the three times are  
 410 showed in the following table.

$t_1$	$t_2$	$t_3$
$\Delta$	$\bullet$	$*$

In Figure 7, we show a solution of type  $m_2 - L_2 - m_2$ . The orbit is obtained  
 starting with initial conditions on  $W^s(L_2^-)$ . On the left, we can see the projec-  
 tion in the synodic configuration space, on the right, the projection in the  $\mathbf{Z}$   
 415 coordinates. Initially, for a range of time including  $t_1$ , the particle surrounds the  
 fixed position of  $m_2$  in synodic coordinates (the particle spins around  $m_2$  on  
 and on along its parabolic orbit in sidereal coordinates). At time  $t_2$  the particle  
 is close to the collinear point  $L_2$  in synodic coordinates, which corresponds to a  
 quasi collinear configuration where the three bodies are almost aligned. As the  
 420 time increases, the particle gets captured around  $m_2$ .

In Figure 8, we show two orbits: one of type  $m_1 - L_2 - m_2$  (orbit “a”) and  
 one of type  $m_1 - L_2 - m_1$  (orbit “b”). In both cases, we have started on  $W^u(L_2^-)$




Figure 7: Solution of type  $m_2 - L_2 - m_2$  of the global system. Left: projection in the synodic configuration space. Right: projection in the inertial system  $\mathbf{Z}$ . The marks with labels  $m_0, m_1, m_2$  denote the position of the primaries and the particle at three different times  $t_1 < t_2 < t_3$ . The position at  $t_2$  corresponds to the passage close to  $L_2$ .

(with  $\theta$  close to  $-\pi/2$ ). We want to stress that whereas along the span of time from capture around  $m_1$  to the close passage to  $L_2$  both orbits are quite close to each other, after the passage close to the collinear configuration, they evolve in a complete different way. One orbit is captured by  $m_2$  whereas the other one is captured by  $m_1$ . So we remark that, some times, a small change in the initial conditions may give rise to rather different behaviors. Of course this is due to the presence of the invariant manifold of  $L_2$ .

In Figure 9 two solutions with close passages to two equilibrium points are shown: one of type  $m_2 - L_3 - L_2 - m_2$  and the other of type  $m_2 - L_3 - L_2 - m_1$ . In this case, the orbits are almost coincident for  $t \leq t_2$ , leaving a neighborhood of  $m_2$  and having a passage close to a collinear configuration. Then, they have different evolutions forwards in time. The invariant manifold of  $L_2^+$  is the responsible for the separation of both trajectories: while one particle goes to collision with  $m_1$ , the other one goes to collision with  $m_2$ . Clearly, in between there must exist a trajectory belonging to  $W^s(L_2^+)$  connecting  $m_2$  with  $L_2^+$ .

Other examples are shown in Figure 10. The left plot shows two orbits visiting three equilibrium points: orbit “a” is a connecting orbit of type  $m_2 - L_2 - L_4 - L_3 - m_2$  whereas orbit “b” is an escape one. It is also clear that there




Figure 8: Solutions of type  $m_1 - L_2 - m_1$  and  $m_1 - L_2 - m_2$ . Top: projection in the synodic configuration space. Bottom: projection in the inertial system  $\mathbf{Z}$ . The marks with labels  $m_0, m_1, m_2$  denote the position of the primaries and the two particles at three different times  $t_1 < t_2 < t_3$ . The position at  $t_2$  corresponds to the passage close to  $L_2$ .

is an intermediate heteroclinic orbit from  $m_2$  to  $L_3^+$  with a close passage  $L_4$ . Similarly the right plot shows two orbits: a solution orbit of type  $m_2 - L_3 - L_5 - m_1$ , an escape one  $m_2 - L_3 - L_5 - \infty$  and there must exist an intermediate one connecting  $m_2$  and  $L_5^+$ .

445 Finally starting with initial conditions on  $W^s(L_4^-)$  or  $W^u(L_4^-)$  and  $\theta$  close to  $-\pi/2$ , we show in Figure 11 two orbits with a close passage to a triangular




Figure 9: Solutions of type  $m_2 - L_3 - L_2 - m_2$  and  $m_2 - L_3 - L_2 - m_1$  of the global system. Top: projection in the synodic configuration space of both trajectories. Bottom: projections in the inertial system  $\mathbf{Z}$ . The marks with labels  $m_0, m_1, m_2$  denote the position of the primaries and the two particles at three different times  $t_1 < t_2 < t_3$ . The position at  $t_2$  correspond to the passage close to  $L_3$ .

configuration: one of type  $m_2 - L_4 - m_1$  (left) and another one of type  $m_2 - L_2 - L_4 - m_2$ .

#### 4.2. Symmetric connecting orbits

450 The procedure to find initial conditions that lead to symmetric connecting orbits is based on symmetries (7) and (8). On one hand, using the symmetry




Figure 10: Projection in the synodic configuration space of solutions of the global system. Left: one solution is of type  $m_2 - L_2 - L_4 - L_3 - m_2$  and the second one is close to the first one but finally escapes. Right: one orbit is of type  $m_2 - L_3 - L_5 - m_1$  while the other escapes.




Figure 11: Projection in the synodic configuration space of solutions of the global system. Left: of type  $m_2 - L_4 - m_1$ . Right: of type  $m_2 - L_2 - L_4 - m_2$ .

(7) if an orbit leaves a neighborhood of one primary, and crosses the section  $\theta = 0$  such that  $y = x' = 0$ , the trajectory will end at the same primary. This is a connection  $m_i - m_i$  symmetric with respect to the  $x$  axis on the  $(x, y)$  plane.

455 On the other hand, using both (7) and (8), if an orbit leaves a neighborhood of one primary, and crosses the section  $\theta = 0$  such that  $x = y' = 0$ , the trajectory will end at the other primary. This is a connection  $m_i - m_j$ ,  $i \neq j$  symmetric with respect to the  $y$  axis on the  $(x, y)$  plane.

We have explored these two kind of connections in the following way. Take

460 initial conditions at  $\theta = 0$  in the plane  $(x, y')$  (connecting orbits for the same primary) and in the plane  $(y, x')$  (connecting orbits between different primaries). Integrate such initial conditions forwards and classify them, using the  $C$ -criterion, depending whether the orbit is a collision orbit with one of the primaries or the orbit escapes to infinity.

465 Notice that in some cases the integration cannot be done until  $C = 8$ , when the  $C$ -criterion is applied, due to the fact that the orbit has a close encounter to one primary. At these points a binary collision regularization should be performed in order to continue the integration. Instead, we classify the orbit as “possible collision with the primary”.

470 To avoid unnecessary computations, we discard initial conditions of the planes  $(x, y')$  and  $(y, x')$  at  $\theta = 0$  such that the value of the Jacobi function  $C$  is greater than  $C_3$  or  $C_2$ , so that the  $C$ -criterion applies directly without integration. More precisely, the curves  $C = C_3$  and  $C = C_2$  determine a subset of the escape and capture regions. Notice that on the plane  $(y, x')$  the curve  
475  $C = C_2$  does not play any role. Moreover, using the symmetries of the problem, it is enough to do the computation in the positive  $x$  and  $y$  planes respectively. The detailed regions of escape and capture on the  $(x, y')$  and  $(y, x')$  planes are shown in Figure 12, left and right, respectively.

As we saw in the previous section, the codimension 1 invariant manifolds  
480 associated to the collinear equilibrium points separate the different types of orbits and these are precisely the boundaries of the escape and capture regions plotted. In order to show this assertion, we consider a region where three different regions (escape, capture around  $m_1$  and capture around  $m_2$ ) meet, for example in the  $(y, x')$  plane, see Figure 13 left. More precisely, we focus on  
485  $[0.52952, 0.52968] \times [-3.33108, -3.33093]$ , around a point where three different regions meet, see Figure 13 right. We take a circle centered at the common point to the three different regions and a small radius, for instance 0.00005. We take initial conditions on this circle parameterized by an angle  $\varphi$  varying from 0 to  $2\pi$  counterclockwise, and consider the corresponding orbits. We show the final  
490 evolution of these orbits varying  $\varphi$  in Figure 14. We start at a given point, la-




Figure 12: Regions of escape (white regions) and capture around  $m_1$  (blue) and  $m_2$  (red) in the  $(x, y')$  (left) and  $(y, x')$  (right) planes. The regions in green and light blue correspond to orbits that reach a neighborhood of  $m_1$  or  $m_2$ , respectively, but the integration has stopped without classifying the point. The curves  $C = C_3$  and  $C = C_2$  correspond to the points at which the Jacobi function values  $C_3$  and  $C_2$  respectively.

beled by “a”. This point belongs to the region of capture orbit around  $m_2$ . As  $\varphi$  increases, the orbits are of the same type until the boundary between this region and the escape region is crossed, see orbits labeled “b” and “c”. We can see in Figure 14 that there is a precise value of  $\varphi$ , such that the orbit tends asymptotically to  $L_3$ . That orbit belongs to  $W^s(L_3^+)$ , which separates the capture orbits around  $m_2$  and escape orbits. Increasing again  $\varphi$ , we obtain escape orbits until the boundary between escape and capture around  $m_1$  is crossed, orbits “d” and “e”. We can see that the point in that boundary corresponds to an orbit that belongs to  $W^s(L_1^+)$ . Finally, increasing  $\varphi$  we have points in the capture region around  $m_1$  until the boundary with the capture region around  $m_2$  is reached again, orbits “f” and “a”. In this case, the boundary corresponds to an orbit on  $W^s(L_2^+)$ . Therefore, the manifolds of  $L_i^+$ ,  $i = 1, 2, 3$  are the boundaries that separate the different behaviors of capture and escape. The initial conditions of the orbits labeled as “a,” . . . , “f” are shown in Table 2. We also want to stress that the common point to the three regions is precisely associated to an orbit tending to  $L_5^+$ . Notice that all the orbits have a close passage to  $L_5$ .




Figure 13: Regions of escape (white regions) and capture around  $m_1$  (blue) and  $m_2$  (red) in the  $(y, x')$  plane. The regions in green and light blue correspond to points that reach a neighborhood of  $m_1$  or  $m_2$ , respectively, but the integration has stopped without classifying the point. The plot on the right is a magnification.




Figure 14: Orbits showing the transitions between capture around  $m_1$ ,  $m_2$  and escape. The separatrix orbits of these transitions are asymptotic orbits tending to  $L_i$ ,  $i = 1, 2, 3$ .

## 5. Conclusions

All possible trajectories connecting A and B (A and B being an equilibrium point, or a collision with a primary, or escape) have been established, except the ones that require a regularization of the singularities of the equations of motion. This was done using the map of heteroclinic connections and the  $C$ -criterion.

A zoo of symmetric and nonsymmetric connecting orbits is presented using different strategies. The first one is based on the knowledge of the invariant

	$y$	$x'$
a	0.5296428592578155	-3.3309921580556328
b	0.5295965978644199	-3.3309718860907380
c	0.5295965977084045	-3.3309718861089928
d	0.5296063350369673	-3.3310713928937119
e	0.5296063351935617	-3.3310713928813760
f	0.5296499612337321	-3.3310060964681765

Table 2: Initial conditions corresponding to the orbits a,...,f shown in Figure 14.

manifolds associated to the equilibrium points while in the second one, the  
515 most important role is played by the symmetries of the problem.

In the case of interacting disk galaxies (each galaxy treated as a point mass  
embedded by a disk of test or massless particles) the present model shows that,  
after a galaxy close encounter, the regions of the phase space where the test par-  
ticles remain or not around each galaxy are confined by the invariant manifolds  
520 of the equilibrium points.

## 6. Acknowledgments

The authors want to thank the fruitful discussions with Dr. Joaquin Delgado  
(UAM-Iztapalapa), who in particular proposed the basic ideas to show the  
heteroclinic connection between  $L_4^+$  and  $L_3^+$ . The authors also want to thank  
525 the conversations with Dr. Martha Álvarez in relation with this problem.

E. Barrabés has been supported by the Spanish grant MTM2013-41168,  
and the Catalan grant 2014 SGR1145. J.M. Cors has been supported by  
MINECO/FEDER grants numbers MTM2008-03437 and MTM2013-40998 and  
by AGAUR grant number 2014 SGR568. M. Ollé has been supported by the  
530 Spanish MINECO/FEDER grant MTM2012-31714 and the Catalan grant 2014  
SGR504.

## References

- [1] M. Alvarez, J.M. Cors, and J. Delgado. On final evolutions in the restricted planar parabolic three-body problem. *Celestial Mech. Dynam. Astronom.*, 95(1-4):173–200, 2006. doi: 10.1007/s10569-006-9018-9. 535
- [2] J.J. Condon, G. Helou, D.B. Sanders, and B.T. Soifer. The "taffy" galaxies ugc 12914/5. *Astronomical Journal*, 105(5):1730–1736, 1993.
- [3] J. M. Cors and J. Llibre. The global flow of the parabolic restricted three-body problem. *Celestial Mech. Dynam. Astronom.*, 90(1-2):13–33, 2004. 540 doi: 10.1007/s10569-004-4917-0.
- [4] J.M.A. Danby. *Fundamentals of Celestial Mechanics (Second Edition)*. Willmann-Bell, Inc., 1992.
- [5] M. Faintich. Application of the restricted hyperbolic three-body problem to a star-sun-comet system. *Celestial Mechanics and Dynamical Astronomy*, 6:22–26, 1972. 545
- [6] M.M. Fragner and R.P. Nelson. Giant planet formation in stellar clusters: The effects of stellar fly-bys. *Astronomy and Astrophysics*, 505(2):873–889, 2009.
- [7] G.I. Günthardt, E.L. Agüero, I. Rodrigues, and R.J. Díaz. Spectroscopic properties and dynamical evolution of the merging system am 1003-435. 550 *Astronomy and Astrophysics*, 453(3):801–808, 2006.
- [8] A. Haro and J.M. Mondelo. Seminumerical algorithms for computing invariant manifolds of vector fields at fixed points. In *The parameterization method for invariant manifolds: from theory to effective computations*. Preprint, 2014. 555
- [9] K. Meyer and Q.D. Wang. Global phase structure of the restricted isosceles three-body problem with positive energy. *Trans. Amer. Math. Soc.*, 338(1): 311–336, 1993. doi: 10.2307/2154458.

- [10] P.M.S. Namboodiri, R.K. Kochhar, and S.M. Alladin. On the formation  
560 of bridges and tails in interacting galaxies. ii. *Bull. Astr. Soc. India*, 15:  
186–192, 1987.
- [11] L.M. Perko. *Differential equations and dynamical systems*. Springer-Verlag,  
New York, second edition, 1996.
- [12] S. Pfalzner, P. Vogel, J. Scharwchter, and C. Olczak. Parameter study of  
565 star-disc encounters. *Astronomy and Astrophysics*, 437(3):967–976, 2005.
- [13] D.G. Saari. *Collisions, rings, and other Newtonian N-body problems*, vol-  
ume 104 of *CBMS Regional Conference Series in Mathematics*. Published  
for the Conference Board of the Mathematical Sciences, Washington, DC;  
by the American Mathematical Society, Providence, RI, 2005.
- 570 [14] M. Steinhausen, C. Olczak, and S. Pfalzner. Disc-mass distribution in  
star-disc encounters. *Astronomy and Astrophysics*, 538, 2012.
- [15] V. Szebehely. *Theory of Orbits. The Restricted Problem of Three Bodies*.  
Academic Press, Inc., 1967.
- [16] A. Toomre and J. Toomre. Galactic bridges and tails. *The Astrophysical*  
575 *Journal*, 178:623–666, 1972.