

Arnold diffusion in a model of dissipative system *

Samuel W. Akingbade[†], Marian Gidea[‡], and Tere M-Seara[§]

Abstract. For a mechanical system consisting of a rotator and a pendulum coupled via a small, time-periodic Hamiltonian perturbation, the Arnold diffusion problem asserts the existence of ‘diffusing orbits’ along which the energy of the rotator grows by an amount independent of the size of the coupling parameter, for all sufficiently small values of the coupling parameter. There is a vast literature on establishing Arnold diffusion for such systems. In this work, we consider the case when an additional, dissipative perturbation is added to the rotator-pendulum system with coupling. Therefore, the system obtained is not symplectic but conformally symplectic. We provide explicit conditions on the dissipation parameter, so that the resulting system still exhibits energy growth. The fact that Arnold diffusion may play a role in systems with small dissipation was conjectured by Chirikov. In this work, the coupling is carefully chosen, however the mechanism we present can be adapted to general couplings and we will deal with the general case in future work.

1. Introduction. The Arnold diffusion problem [Arn64] broadly refers to a universal mechanism of instability for multi-dimensional Hamiltonian systems that are small perturbations of integrable ones. Through this mechanism, chaotic transfers of energy take place between subsystems of a given Hamiltonian system, which, in particular, can lead to significant growth of energy of one of the subsystems over time. Chirikov [Chi79] conjectured that Arnold diffusion may play a role in systems with small dissipation as well.

Studying Hamiltonian systems with small dissipation is important for applications, as many real-life physical systems experience some energy loss over time.

A significant class of examples is furnished by Celestial Mechanics, on the motion of celestial bodies under mutual gravity. As the gravitational force is conservative, such systems are usually modeled as Hamiltonian systems. Nevertheless dissipative forces are present in real-world systems, including tidal forces, Stokes drag, Poynting-Robertson effect, Yarkowski/Y-ORP effects, atmospheric drag, and their effect may accumulate in the long run. While some of these effects may be negligible over relatively short time scales, others, for instance Earth’s atmospheric drag on artificial satellites, can have significant effects over practical time scales. See, e.g. [MNF87, Cel07, RR17].

Another class of examples is given by energy harvesting devices. Some of these devices consist of systems of oscillating beams made of piezoelectric materials, where on the one hand there is dissipation due to mechanical friction, and on the other hand there is external

*Submitted to the editors DATE. S.W.A. and M.G. were partially supported by NSF grant DMS-1814543. The work of T.M-S. was partially supported by the grants PGC2018-098676-B-I00 and PID-2021-122954NB-I00 funded by MCIN/AEI/10.13039/501100011033 and “ERDF A way of making Europe”. This work is also supported by the Spanish State Research Agency, through the Severo Ochoa and María de Maeztu Program for Centers and Units of Excellence in R&D (CEX2020-001084-M). T.M-S. was partially supported by the Catalan Institution for Research and Advanced Studies via an ICREA Academia Prize 2019.

[†]Yeshiva University, Department of Mathematical Sciences, New York, NY 10016, USA (sakingba@mail.yu.edu)

[‡]Yeshiva University, Department of Mathematical Sciences, New York, NY 10016, USA (Marian.Gidea@yu.edu).

[§]Departament de Matemàtiques and IMTECH, Universitat Politècnica de Catalunya (UPC), Diagonal 647, 08028 Barcelona, Spain, Centre de Recerca Matemàtica, Barcelona, Spain. (Tere.M-Seara@upc.edu)

forcing, owed to the movement of the device, that triggers the beams to oscillate. See, e.g. [MH79, EHI09, Gra17].

Of course, there are many other examples. In this paper we consider a simple model of a mechanical system, consisting of a rotator and a pendulum with a small, periodic coupling, subject to a small dissipative perturbation. Coupled rotator-pendulum systems are fundamental models in the study of Arnold diffusion in Hamiltonian systems. Adding a dissipative perturbation results in a system that is non-Hamiltonian. The symplectic structure changes into a conformally symplectic one. We show that such a system exhibits Arnold diffusion, in the sense that there exist pseudo-orbits for which the energy of the rotator subsystem grows by some quantity that is independent of the smallness parameter. (By a pseudo-orbit here we mean a sequence of orbit segments of the flow such that the endpoint of each orbit segment is ‘close’ to the starting point of the next orbit segment in the sequence.) We note that for the unperturbed rotator-pendulum system, the energy of the rotator subsystem is conserved. The small, periodic coupling added to the system makes the rotator undergo small oscillations in energy, while the dissipative perturbation typically yields a loss in energy. The physical significance of our result is that, despite the dissipation effects, it is possible to overall gain a significant amount of energy over time.

Specifically, the unperturbed rotator-pendulum system is given by a Hamiltonian of the form

$$H_0(p, q, I, \theta) = h_0(I) + h_1(p, q),$$

with $z = (p, q, I, \theta) \in \mathbb{R} \times \mathbb{T}^1 \times \mathbb{R} \times \mathbb{T}^1$, where $h_0(I)$ represents the Hamiltonian of the rotator, and h_1 represents the Hamiltonian of the pendulum, and $\mathbb{T}^1 = \mathbb{R}/2\pi\mathbb{Z}$. The perturbed system is of the form

$$(1.1) \quad \dot{z} = J\nabla_z H_0(z) + \varepsilon J\nabla_z H_1(z, t) + \mathcal{X}_\lambda(z),$$

where $H_1(z, t)$ is a Hamiltonian that is 2π -periodic in time t , $\varepsilon \geq 0$ is the size of the coupling, $\mathcal{X}_\lambda(z)$ is a dissipative vector field depending on some dissipation parameter $\lambda = \lambda(\varepsilon) > 0$, and

$$J = \begin{pmatrix} J_2 & 0 \\ 0 & J_2 \end{pmatrix} \text{ where } J_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Technical conditions on $h_0, h_1, H_1, \mathcal{X}_\lambda$ will be given in Section 3. Under those conditions, the phase space of the perturbed system has a 3-dimensional Normally Hyperbolic Invariant Manifold (NHIM from now on), which contains a 2-dimensional invariant torus that is an attractor for the dynamics in the NHIM. This torus creates a ‘barrier’ for the existence of diffusing orbits by using only the ‘inner dynamics’ (i.e., the dynamics restricted to the NHIM). The main question is whether there are diffusing orbits crossing this ‘barrier’ by combining the ‘inner dynamics’ with the ‘outer dynamics’ (i.e., the dynamics along homoclinic orbits to the NHIM).

We show that there exist $C > 0$ and $\varepsilon_0 > 0$ such that, for all $0 < |\varepsilon| < \varepsilon_0$, there exists a pseudo-orbit $z(t)$, $t \in [0, T]$, of (1.1), such that

$$I(T) - I(0) > C \text{ for some } T > 0.$$

More technical details will be given in Theorem 4.1. In order for the above result to be of practical interest, the above solution $z(t)$ should be chosen such that at the beginning $(I(0), \theta(0))$ is below (relative to I) the aforementioned attractor, and at the end $(I(T), \theta(T))$ is above the aforementioned attractor. Indeed, it is possible to increase I by starting below the attractor and moving towards the attractor under the effect of the dissipation alone; obviously, such a solution is not of practical interest.

2. Conservative vs. dissipative systems. Arnold's conjecture on Hamiltonian instability originated with an example of a rotator-pendulum system with a small, time-periodic Hamiltonian coupling of special type [Arn64]. In his example, in the absence of the coupling, the phase space of the rotator forms a normally hyperbolic invariant manifold (NHIM) foliated by 'whiskered', rotational tori, which have stable and unstable invariant manifolds that coincide. The coupling in Arnold's example was specially chosen so that it vanishes on the family of invariant tori, and so the tori are preserved. These tori constitute 'barriers' for the existence of diffusing orbits, since orbits in the NHIM always move along these tori and thus cannot increase their action variable. At the same time the coupling splits the stable and unstable manifolds, so that the unstable manifold of each torus intersects transversally the stable manifolds of nearby tori. Thus, one can form 'transition chains' of tori, and show that, by interspersing the 'outer dynamics' along the homoclinic orbits to the NHIM with the 'inner dynamics' along the tori, one can obtain 'diffusing' orbits along which the energy of the rotator exhibits a significant growth. Arnold conjectured that this mechanism of diffusion occurs in close to integrable general systems.

However, in the case of a general coupling not all of the invariant tori in the NHIM are preserved. The KAM theorem yields a Cantor set of tori that survive from the unperturbed case, with gaps in between. The splitting of the stable and unstable manifold makes the unstable manifold of each torus intersect transversally the stable manifolds of sufficiently close tori, however the size of the splitting is in general smaller than the size of the gaps between tori. This is known as the 'large gap problem'. It was overcome, for instance, by forming transition chains that, besides rotational tori, also include 'secondary' tori created by the perturbation [DdlS00, DLS06]. Other geometric mechanisms use transition chains that include, besides rotational tori, Aubry-Mather sets [GR13]. Subsequently, [GdlMS20] described a general mechanism of diffusion that relies mostly on the outer dynamics, and uses only the Poincaré recurrence of the inner dynamics (which is automatically satisfied in Hamiltonian systems over regions of bounded measure).

The references mentioned above encompass geometric ideas that we can adapt to the dissipative case. However, there are many other geometric mechanisms that have been used in the Arnold diffusion problem, such as those in [CG94, BT99, DdlS00, Tre02, Tre04, DdlS06a, DdlS06b, Pif06, GT08, DH09, Tre12, GdlL17, GT17a, GM22]. A variational program for the Arnold diffusion was formulated in [Mat04, Mat12] for systems close to integrable. Global variational methods for diffusion have been used in this setting for convex Hamiltonians [CY09, KZ15, BKZ16, CX19, KZ20]. A hybrid program combining geometric and variational methods was started in [BB02, BBB03].

The case of a rotator-pendulum system subject to a non-Hamiltonian perturbation (consisting of time-periodic Hamiltonian coupling and a dissipative force) which we consider in this

paper, has very different geometric features from the conservative case. The dissipation added to the Hamiltonian system is a singular perturbation – the system with positive dissipation leads to attractors inside the NHIM, which can contain at most one invariant torus. Poincaré recurrence does not hold for dissipative systems. The stable and unstable manifolds of the NHIM do not necessarily intersect. Therefore, the mechanism used for proving diffusion in the Hamiltonian case does not carry over to the non-Hamiltonian case.

To provide some intuition, we illustrate on a couple of basic examples some possible effects of dissipation on the geometry of Hamiltonian systems.

Example 2.1. The first example is the standard map.

The (conservative) standard map, which can be viewed as the time-one map of a non-autonomous Hamiltonian system representing a ‘kicked rotator’, is given by

$$(2.1) \quad \begin{aligned} I' &= I + \varepsilon \sin(\theta), \\ \theta' &= \theta + I + \varepsilon \sin(\theta), \end{aligned}$$

where ε is the perturbative parameter, and I, θ are defined $(\text{mod } 2\pi)$. This is a symplectic twist map; the symplecticity condition being $dI' \wedge d\theta' = dI \wedge d\theta$ and the twist condition being $\frac{\partial \theta'}{\partial I} \neq 0$. When $\varepsilon = 0$ the resulting map is the time-one map of a rotator, and is given by

$$(2.2) \quad \begin{aligned} I' &= I, \\ \theta' &= \theta + I. \end{aligned}$$

It is an integrable twist map, with all level sets of I being rotational invariant circles on which the motion is a rigid rotation of frequency $\omega(I) = I$. For $0 < \varepsilon \ll 1$, the KAM theorem asserts that there is a positive measure set of invariant circles, of Diophantine frequencies, which survive the perturbation. The measure of the set of the KAM circles tends to 1 as $\varepsilon \rightarrow 0$. On the other hand, when $\varepsilon > 0$ increases, fewer and fewer invariant circles survive, and eventually only one invariant circle is left. The last rotational invariant circle for the standard map has frequency $\omega = \frac{1+\sqrt{5}}{2}$, which is the golden mean [Gre79]. See Fig. 1a.

The dissipative standard map is defined as

$$(2.3) \quad \begin{aligned} I' &= (1 - \lambda)I + \mu + \varepsilon \sin(\theta), \\ \theta' &= \theta + (1 - \lambda)I + \mu + \varepsilon \sin(\theta), \end{aligned}$$

where λ is the dissipative parameter, $0 \leq \lambda < 1$, and μ is the drift parameter; $\lambda = 0$ corresponds to no dissipation. The map is no longer symplectic, but conformally symplectic, that is $dI' \wedge d\theta' = (1 - \lambda)dI \wedge d\theta$, and still satisfies a twist condition.

When $\varepsilon = 0$, the resulting map

$$(2.4) \quad \begin{aligned} I' &= (1 - \lambda)I + \mu, \\ \theta' &= \theta + (1 - \lambda)I + \mu, \end{aligned}$$

has a single rotational invariant circle $I = \frac{\mu}{\lambda}$ of frequency $\omega_* := \frac{\mu}{\lambda}$. The KAM theorem for conformally symplectic systems asserts that for each $0 < \varepsilon \ll 1$ there is one rotational invariant circle, of Diophantine frequency, that survives the perturbation, and that circle is a

local attractor for the system (see, e.g. [CC09, CCdIL13b, CCDIL13a, CCdIL20]). In order for the surviving circle to be of Diophantine frequency ω_* , we need to properly adjust the drift parameter $\mu = \mu(\omega_*, \varepsilon)$. See Fig. 1b and Fig. 1c.

We can rewrite the dissipative standard map in terms of a frequency parameter ω_* rather than in terms of the drift parameter $\mu = \lambda\omega_*$ obtaining:

$$(2.5) \quad \begin{aligned} I' &= I - \lambda(I - \omega_*) + \varepsilon \sin(\theta), \\ \theta' &= \theta + I - \lambda(I - \omega_*) + \varepsilon \sin(\theta). \end{aligned}$$

In this case, by the persistence of normal hyperbolicity of the torus given by $I = \omega_*$, as $0 < \varepsilon \ll 1$ is varied, there exists an invariant torus of frequency $\omega = \omega(\varepsilon)$ close to ω_* ; not all frequencies ω yield KAM circles but only those $\omega(\varepsilon)$ which are Diophantine.

Example 2.2. The second example is the pendulum, given by the Hamiltonian

$$h_1(p, q) = \frac{p^2}{2} + (\cos(q) - 1).$$

As it is well known, the pendulum has a hyperbolic fixed point whose stable and unstable manifolds coincide (see Fig. 2a).

When dissipation is added to the pendulum

$$\begin{aligned} \dot{p} &= -\lambda p + \sin(q), \\ \dot{q} &= p, \end{aligned}$$

the origin is again a hyperbolic fix point with eigenvalues $\pm\sqrt{1 + (\frac{\lambda}{2})^2} - \frac{\lambda}{2}$. Nevertheless, its stable and unstable manifolds cease to intersect, for dissipative coefficient $\lambda > 0$ (see Fig. 2b).

However, when both dissipation and periodic forcing are added to the pendulum,

$$\begin{aligned} \dot{p} &= -\lambda p + \sin(q) + \varepsilon \sin(t), \\ \dot{q} &= p, \end{aligned}$$


for certain parameter values $\lambda > 0$ and $\varepsilon > 0$, the time- 2π map exhibits chaotic attractors in the Poincaré section (see Fig. 2c).

These simple examples illustrate that adding dissipation to a Hamiltonian system typically destroys – sometimes dramatically – some of the geometric structures – KAM tori, homoclinic connections – that are relevant in Arnold’s mechanism of diffusion, and creates new geometric structures – attractors – that act as barriers for diffusion. On the other hand, the addition of forcing can compensate the effects of dissipation.


3. Model. The model that we consider is described by an integrable Hamiltonian system subject to a time-dependent, Hamiltonian perturbation (or coupling), and to a second, non-Hamiltonian, perturbation that is dissipative.

The unperturbed Hamiltonian corresponds to an uncoupled rotator-pendulum system and is given by


$$(3.1) \quad H_0(p, q, I, \theta) = \frac{I^2}{2} + \frac{p^2}{2} + (\cos q - 1),$$




(a) Conservative standard map



(b) Dissipative standard map with unadjusted drift



(c) Dissipative standard map with adjusted drift



(d) The basins of attraction of the attractors appearing in (c), using a colour scale based on rotation numbers (Credit Matteo Manzi)

Figure 1. *The conservative and dissipative standard map*

where $(p, q, I, \theta) \in \mathbb{R} \times \mathbb{T}^1 \times \mathbb{R} \times \mathbb{T}^1$, which is endowed with the standard symplectic structure $\omega = dp \wedge dq + dI \wedge d\theta$.

For the rotator part of the Hamiltonian, given by $h_0(I) = \frac{I^2}{2}$, each level set $I = \text{constant}$ is invariant under the flow of h_0 , and the corresponding dynamics is a rigid rotation of frequency

$$(3.2) \quad \omega(I) := \frac{\partial h_0}{\partial I} = I.$$

The pendulum part of the Hamiltonian, is given by

$$(3.3) \quad h_1(p, q) = \frac{p^2}{2} + (\cos q - 1),$$




Figure 2. The pendulum

183 it has a hyperbolic fixed point at $(p, q) = (0, 0)$ and an elliptic fixed point at $(p, q) = (0, \pi)$.
 184 The stable and unstable manifolds of the hyperbolic fixed point $(0, 0)$ coincide, and can be
 185 parametrized as

$$186 \quad (3.4) \quad (p_0(t), q_0(t)) = \left(\pm \frac{2}{\cosh t}, 4 \arctan e^{\pm t} \right).$$

187 Since the system H_0 is uncoupled, I is a conserved quantity and so each hypersurface
 188 $\{I = \text{const.}\}$ constitutes a barrier for the dynamics of H_0 : there are no trajectories along
 189 which the variable I can change.

190 When we add the time-dependent, Hamiltonian perturbation, we have

$$191 \quad (3.5) \quad H_\varepsilon(p, q, I, \theta, t) = H_0(p, q, I, \theta) + \varepsilon H_1(p, q, I, \theta, t),$$

192 where $t \in \mathbb{T}^1$, meaning that the perturbation H_1 is 2π -periodic in time.

193 We will assume that H_1 is of the form

$$194 \quad (3.6) \quad H_1(p, q, I, \theta, t) = f(q) \cdot g(\theta, t).$$

195 The dissipative perturbation is given by a vector field \mathcal{X}_λ that is added to the Hamiltonian
 196 vector field $J\nabla H_\varepsilon$ of H_ε , where

$$197 \quad (3.7) \quad \mathcal{X}_\lambda(p, q, I, \theta) = (-\lambda p, 0, -\lambda(I - \omega_*), 0),$$

198 where λ is the dissipation coefficient, and ω_* is a fixed Diophantine frequency. For the moment,
 199 we will treat λ as an independent parameter, but for most of the paper we will consider λ of
 200 the form $\lambda = \varepsilon \rho$, with ρ being a sufficiently small independent parameter. In our main result
 201 Theorem 4.1 we will use $\lambda = \varepsilon \rho(\varepsilon)$ where $0 < \rho(\varepsilon) = \frac{\bar{\rho}}{\log \frac{1}{\varepsilon}} \ll 1$, where $\bar{\rho}$ is a positive constant.

The system of interest is

$$(3.8) \quad \dot{z} = J\nabla_z H_0(z) + \varepsilon J\nabla_z H_1(z, t) + \mathcal{X}_\lambda(z), \quad z = (p, q, I, \theta).$$

We obtain the following equations:

$$(3.9) \quad \begin{cases} \dot{p} = \sin(q) - \varepsilon f'(q) \cdot g(\theta, t) - \lambda p, \\ \dot{q} = p, \\ \dot{I} = -\lambda(I - \omega_*) - \varepsilon f(q) \cdot \frac{\partial g}{\partial \theta}(\theta, t), \\ \dot{\theta} = I. \end{cases}$$

As we shall see, the dissipative perturbation yields the existence of attractors for the dynamics restricted to the NHIM, that is, the dynamics in the (I, θ) -variables. In particular, we can have attractors that act as barriers on the NHIM, in the sense that they separate the NHIM into topologically non-trivial connected components. As all trajectories within the basin of attractors move towards the attractors, the action I will increase along some trajectories and will decrease along some other trajectories, however there are no trajectories within the NHIM that start on one side of the attractor and end on the other side.

Below, we will consider two concrete examples of Hamiltonian perturbations:

Vanishing perturbation: H_1 vanishes at $(p, q) = (0, 0)$

$$(3.10) \quad \begin{aligned} f(q) &= \cos(q) - 1, \\ g(\theta, t) &= a_{00} + a_{10} \cos \theta + a_{01} \cos t. \end{aligned}$$

Non-vanishing perturbation: H_1 does not vanish at $(p, q) = (0, 0)$

$$(3.11) \quad \begin{aligned} f(q) &= \cos(q), \\ g(\theta, t) &= a_{00} + a_{10} \cos \theta + a_{01} \cos t. \end{aligned}$$

Above, a_{00} , a_{10} , and a_{01} are real numbers with $a_{10}a_{01} \neq 0$.

Remark 3.1. The choice of the coupling of the form $H_1(p, q, I, \theta, t) = f(q)g(\theta, t)$ has been made in order to deal with a simple model. The fact that the function f satisfies $f'(0) = 0$ implies that the normally hyperbolic invariant manifold, which is exhibited by the unperturbed system, is not affected by the perturbation; see Section 5.4. We do not need to invoke the theory of persistence of normally hyperbolic invariant manifolds under perturbation. The function $g(\theta, t)$ can be viewed as a truncation to the first two harmonics of the Fourier expansion of an analytic function. We will deal with the general case with infinitely many harmonics, as well as with perturbations that do not preserve the NHIM, in future work.

Remark 3.2. We note that instead of (3.7) we can consider more general perturbations of the form

$$\mathcal{X}_\lambda(p, q, I, \theta) = (-\lambda_1 p, 0, -\lambda_2(I - \omega_*), 0),$$

for $\lambda_1 = \varepsilon \bar{\rho}_1$ and $\lambda_2 = \varepsilon \frac{\bar{\rho}_2}{\log \frac{1}{\varepsilon}}$ with $\bar{\rho}_1, \bar{\rho}_2 > 0$. One will be able to see that the arguments below also apply to this case and therefore, the main result, stated below, remains valid.

4. Main result.

Theorem 4.1. *Consider the Hamiltonian system (3.1) subject to the time-periodic Hamiltonian perturbation (3.10) or (3.11), and to the dissipative perturbation (3.7), with dissipation coefficient $\lambda = \varepsilon \rho(\varepsilon) = \varepsilon \frac{\bar{\rho}}{\log(\frac{1}{\varepsilon})}$ with $\bar{\rho} > 0$ suitably small.*

Then there exist $0 < I_1 < I_2$ and $\varepsilon_0 > 0$ such that, for every ω_ Diophantine number with $0 < I_1 < \omega_* < I_2$, and every $0 < \varepsilon < \varepsilon_0$, there exist pseudo-orbits $z(t)$, $t \in [0, T]$, such that*

$$I(z(0)) < I_1 \text{ and } I(z(T)) > I_2.$$

Here by a pseudo-orbit $z(t)$ we mean a finite collection of trajectories $z^i(t)$, $t \in [t_i, t_{i+1}]$ of (3.8), for some times $0 = t_0 < t_1 < \dots < t_m = T$, where $m > 0$, such that

$$I(z^0(0)) < I_1 \text{ and } I(z^m(T)) > I_2,$$

$$d(z^i(t_{i+1}), z^{i+1}(t_{i+1})) < \delta(\varepsilon), \text{ for } i = 0, \dots, m-1,$$

for some $\delta(\varepsilon) = O(\varepsilon) > 0$.

The diffusion time along the pseudo-orbit $z(t)$, $t \in [0, T]$, is $T = O(\frac{1}{\varepsilon} \log(\frac{1}{\varepsilon}))$.

Above, we used the notation $f = O(g) = O_{\mathcal{C}^k}(g)$ for a pair of functions f, g satisfying $\|f\|_{\mathcal{C}^k} \leq M \|g\|_{\mathcal{C}^k}$ for some $M > 0$, where $\|\cdot\|_{\mathcal{C}^k}$ is the \mathcal{C}^k -norm for some suitable $k \geq 0$.

We illustrate the phenomenon described by Theorem 4.1 in Fig. 3a, Fig. 3b, Fig. 3c, Fig. 3d. In Fig. 3a, using the inner dynamics alone, orbits with $I(0) < \omega_*$ cannot pass beyond the attractor shown in Fig. 3b. However, using both the inner and outer dynamics, there are orbits with $I(0) < \omega_*$ that end up with $I(T) > \omega_*$, as shown in Fig. 3d. These orbits move close to the separatrix of the pendulum $h_1(q, p) = 0$, as shown in Fig. 3c.

Theorem 4.1 gives us diffusing pseudo-orbits. Applying a Shadowing Lemma type of results similar to those in [Zgl09, GT17b, GdlLMS20, CG], we will be able to show that there exist true orbits $z(t)$ of (1.1) such that $I(z(0)) < I_1$ and $I(z(T)) > I_2$. We leave the technical details for a future work.

The above pseudo-orbits are such that the end-point of one is $\delta(\varepsilon)$ -close to the starting-point of the next one, where $\delta(\varepsilon) = O(\varepsilon)$. We remark here that we can also obtain pseudo-orbits with $\delta(\varepsilon) = O(\varepsilon^p)$, for any $p \geq 1$, with the same diffusion time order $T = O(\frac{1}{\varepsilon} \log(\frac{1}{\varepsilon}))$.

In practical applications one can pass from pseudo-orbits to true orbits by applying small controls; for example, in the case of artificial satellites perturbed by atmospheric drag, the small controls can be satellite maneuvers.

Remark 4.2. In Theorem 4.1, the action levels I_1 and I_2 can be chosen explicitly, depending on the Hamiltonian perturbation (3.10) or (3.11) that is considered. See Section 7.7.

The condition on choosing ω_* a Diophantine number between I_1 and I_2 is not necessary for the proof of the theorem; see Section 8.2. The reason for requiring this condition is to be able to apply the KAM theorem for conformally symplectic systems [CCdlL20], which implies the existence of a KAM torus that is an attractor for the inner dynamics, and hence represents a barrier for the inner dynamics. In other words, we want to show that diffusing pseudo-orbits exist even if there is a barrier inside the NHIM.

The choice of the dissipation coefficient $\lambda = \varepsilon \frac{\bar{\rho}}{\log(\frac{1}{\varepsilon})}$ is related to the time $T_h = O(\log(\frac{1}{\varepsilon}))$ required for a point starting in an ε -neighborhood of the NHIM to travel along a homoclinic




Figure 3. *Orbits of the rotator-pendulum system*

orbit and arrive in an ε -neighborhood of the NHIM. This choice of λ implies that $\lambda \cdot T_h = O(\bar{\rho}\varepsilon)$, while the order of the change in action by the scattering map is $O(\varepsilon)$. By choosing a suitable small enough constant $\bar{\rho}$, we can ensure that, when the growth in I by the scattering map competes with the decay in I by the dissipation, which is of order $O(\lambda T_h) = O(\bar{\rho}\varepsilon)$, we will make the former to win against the latter.

If we do not impose that the homoclinic orbits get ε -close to the NHIM, then we can choose a shorter time T_h along the homoclinic orbits and implicitly a larger λ , as long as $\lambda \cdot T_h$ is $O(\varepsilon)$; for example, we can choose $T_h = O(1)$ and $\lambda = \bar{\rho}\varepsilon$ for some $\bar{\rho} > 0$ suitably small.

5. Preliminaries.

5.1. Extended system. Since the perturbation H_1 is time-dependent, it is convenient to consider time as an independent variable t and to work in the extended phase space $\tilde{z} =$

279 $(p, q, I, \theta, s) \in \mathbb{R} \times \mathbb{T}^1 \times \mathbb{R} \times \mathbb{T}^1 \times \mathbb{T}^1$, adding the equation $\dot{s} = 1$ to system (3.9) to obtain:

$$280 \quad (5.1) \quad \begin{aligned} \dot{z} &= J\nabla_z H_0(z) + \varepsilon J\nabla_z H_1(z, s) + \mathcal{X}_\lambda(z), & z &= (p, q, I, \theta) \\ \dot{s} &= 1 \end{aligned}$$

281 We denote by $\tilde{\Phi}_0^t$ the unperturbed extended flow, and by $\tilde{\Phi}_\varepsilon^t$ the perturbed extended flow.

282 **5.2. Normally hyperbolic invariant manifolds.** We briefly recall the notion of a normally
283 hyperbolic invariant manifold (NHIM) [Fen74, HPS77].

284 Let M be a \mathcal{C}^r -smooth manifold, Φ^t a \mathcal{C}^r -flow on M . A submanifold (with or without
285 boundary) Λ of M is a normally hyperbolic invariant manifold (NHIM) for Φ^t if it is invariant
286 under Φ^t , and there exists a splitting of the tangent bundle of TM into sub-bundles over Λ

$$287 \quad (5.2) \quad T_z M = E_z^u \oplus E_z^s \oplus T_z \Lambda, \quad \forall z \in \Lambda$$

288 that are invariant under $D\Phi^t$ for all $t \in \mathbb{R}$, and there exist rates

$$289 \quad \lambda_- \leq \lambda_+ < \lambda_c < 0 < \mu_c < \mu_- \leq \mu_+$$

290 and a constant $C > 0$, such that for all $x \in \Lambda$ we have

$$291 \quad (5.3) \quad \begin{aligned} Ce^{t\lambda_-} \|v\| &\leq \|D\Phi^t(z)(v)\| \leq Ce^{t\lambda_+} \|v\| \text{ for all } t \geq 0, \text{ if and only if } v \in E_z^s, \\ Ce^{t\mu_+} \|v\| &\leq \|D\Phi^t(z)(v)\| \leq Ce^{t\mu_-} \|v\| \text{ for all } t \leq 0, \text{ if and only if } v \in E_z^u, \\ Ce^{|t|\lambda_c} \|v\| &\leq \|D\Phi^t(z)(v)\| \leq Ce^{|t|\mu_c} \|v\| \text{ for all } t \in \mathbb{R}, \text{ if and only if } v \in T_z \Lambda. \end{aligned}$$

292 It is known that Λ is \mathcal{C}^ℓ -differentiable, with $\ell \leq r - 1$, provided that

$$293 \quad (5.4) \quad \begin{aligned} \ell\mu_c + \lambda_+ &< 0, \\ \ell\lambda_c + \mu_- &> 0. \end{aligned}$$

The manifold Λ has associated unstable and stable manifolds, denoted $W^u(\Lambda)$ and $W^s(\Lambda)$, which are tangent to E_Λ^u and E_Λ^s respectively, and $\mathcal{C}^{\ell-1}$ -differentiable. They are foliated by 1-dimensional unstable and stable manifolds (fibers) of points, $W^u(z)$, $W^s(z)$, $z \in \Lambda$, respectively, which are as smooth as the flow, i.e., \mathcal{C}^r -differentiable. These fibers are equivariant in the sense that

$$\begin{aligned} \Phi^t(W^u(z)) &= W^u(\Phi^t(z)), \\ \Phi^t(W^s(z)) &= W^s(\Phi^t(z)). \end{aligned}$$

294 **5.3. The NHIM of the unperturbed system.** We now describe the geometric structures
295 for the unperturbed system corresponding to $\varepsilon = 0$ and $\lambda = 0$. Fix $0 < I_1 < I_2$.

296 The unperturbed system H_0 has a NHIM:

$$297 \quad \Lambda_0 = \{(0, 0, I, \theta) \mid I \in [I_1, I_2], \theta \in \mathbb{T}^1\}.$$

298 The flow restricted to Λ_0 corresponds to the equations of the rotator subsystem:

$$299 \quad (5.5) \quad \begin{cases} \dot{I} = 0 \\ \dot{\theta} = I \end{cases}$$

Hence every level set $I = \text{const.}$ is invariant under the flow. The stable and unstable manifolds of the NHIM Λ_0 coincide, that is

$$W^s(\Lambda_0) = W^u(\Lambda_0) = \{(p, q, I, \theta) \mid h_1(p, q) = 0, I \in [I_1, I_2], \theta \in \mathbb{T}^1\}.$$

where h_1 is the Hamiltonian of the pendulum in (3.3). The contraction/expansion rates along E^s and E^u are ∓ 1 , respectively. For the time- 2π map, the corresponding contraction/expansion rates are $e^{\mp 2\pi}$.

In the extended phase space, we have that $\tilde{\Lambda}_0 = \Lambda_0 \times \mathbb{T}^1$ is a NHIM, and

$$W^s(\tilde{\Lambda}_0) = W^s(\Lambda_0) \times \mathbb{T}^1 = W^u(\Lambda_0) \times \mathbb{T}^1 = W^u(\tilde{\Lambda}_0)$$

for system (5.1).

Note that Λ_0 is also the NHIM for the time- 2π map f_0 of the extended flow $\tilde{\Phi}_0^t$, which represents the first-return map to the Poincaré section

$$\Sigma = \{(p, q, I, \theta, s) \mid s = 0\}.$$

5.4. The inner map of the unperturbed system. Now, let us consider the time- 2π map for the Hamiltonian flow of the rotator:

$$\begin{cases} \dot{I} = 0, \\ \dot{\theta} = I. \end{cases}$$

Solving, we have $I(t) = I_0$ and $\theta(t) = \theta_0 + I_0 t$, which gives the time- 2π -map f_0 :

$$(5.6) \quad f_0(I, \theta) = (I', \theta') = (I, \theta + 2\pi I).$$

Note that f_0 satisfies the twist condition

$$(5.7) \quad \frac{\partial \theta'}{\partial I} = 2\pi > 0.$$

5.5. The model with small dissipation. From now on we will work with small dissipation. We will assume

$$(5.8) \quad \lambda = \varepsilon \rho,$$

where ρ is a free parameter. Consequently, the vector field (3.8) can be written as

$$(5.9) \quad \dot{z} = \mathcal{X}^0(z) + \varepsilon \mathcal{X}^1(z, t; \rho)$$

with $\mathcal{X}^0(z) = J \nabla H_0(z)$ is the unperturbed system (3.1), and

$$(5.10) \quad \mathcal{X}^1(z, t; \rho) = J \nabla H_1(z, t) + \mathcal{X}_\rho,$$

with H_1 given in (3.6) and \mathcal{X}_ρ given in (3.7). Even when we use λ in the notation, we always assume that $\lambda = \varepsilon \rho$.

5.6. The NHIM in the case of vanishing perturbation. In the case when the perturbation H_1 is of the form

$$H_1(p, q, I, \theta, s) = (\cos q - 1) \cdot g(\theta, s),$$

then H_1 vanishes at $(p, q) = (0, 0)$. When εH_1 is added to H_0 , the NHIM $\tilde{\Lambda}_0$ persists as $\tilde{\Lambda}_\varepsilon = \tilde{\Lambda}_0$ for the perturbed system for $\varepsilon > 0$, and the flow restricted to the NHIM is given by (5.5). Consequently, each level set $\{I = \text{constant}\}$ in the NHIM persists.

When we add the dissipation \mathcal{X}_λ , where $\lambda = \varepsilon\rho$, since the (p, q) -components of \mathcal{X}_λ vanish at $(p, q) = (0, 0)$, then the NHIM survives for the perturbed system (3.8) as $\tilde{\Lambda}_\varepsilon = \tilde{\Lambda}_0$ (if we consider λ as an independent parameter, then the perturbed NHIM in general depends on both ε and λ). The induced dynamics on $\tilde{\Lambda}_\varepsilon = \tilde{\Lambda}_0$ is given by

$$(5.11) \quad \begin{cases} \dot{I} = -\lambda(I - \omega_*) \\ \dot{\theta} = I \\ \dot{s} = 1 \end{cases}$$

Note that $\dot{I} = 0 \Leftrightarrow I = \omega_*$. It follows that

$$(5.12) \quad \tilde{A}_\varepsilon = \{(I, \theta, s), I = \omega_*, (\theta, s) \in \mathbb{T}^2\} \subset \tilde{\Lambda}_\varepsilon$$

is a 2-dimensional torus invariant under the flow restricted to $\tilde{\Lambda}_\varepsilon$. This is the only invariant torus for the flow on $\tilde{\Lambda}_\varepsilon$. On \tilde{A}_ε we have $\dot{\theta} = I = \omega_*$ and $\dot{s} = 1$, so the flow along this level set is a linear flow with frequency vector $(\omega_*, 1)$.

By integration of (5.11), we obtain the general solution with initial condition (I_0, θ_0, s_0) as:

$$(5.13) \quad \begin{aligned} I(t) &= (I_0 - \omega_*)e^{-\lambda t} + \omega_*, \\ \theta(t) &= \theta_0 + \frac{1}{\lambda}(I_0 - \omega_*)(1 - e^{-\lambda t}) + \omega_*t, \\ s(t) &= s_0 + t. \end{aligned}$$

Using these explicit formulas, one can see that given $(\omega_*, \theta_0, s_0) \in \tilde{A}_\varepsilon$, if we consider $(I, \theta, s) \in \tilde{\Lambda}_\varepsilon$, where $s = s_0$, $\theta = \theta_0 - \frac{1}{\lambda}(I - \omega_*)$, then

$$(5.14) \quad \|\tilde{\Phi}_\varepsilon^t(I, \theta, s_0) - \tilde{\Phi}_\varepsilon^t(\omega_*, \theta_0, s_0)\| \leq \left(1 + \frac{1}{\lambda^2}\right)^{1/2} |I - \omega_*| e^{-\lambda t} \rightarrow 0 \text{ as } t \rightarrow \infty,$$

showing that \tilde{A}_ε is a global attractor for the flow on $\tilde{\Lambda}_\varepsilon$. (Above we also denoted by $\tilde{\Phi}_\varepsilon^t$ the flow restricted to $\tilde{\Lambda}_\varepsilon$.)

5.7. The inner map in the case of vanishing perturbation. From the explicit solutions of $I(t)$ and $\theta(t)$ in (5.13) with $t = 2\pi$, we have that

$$(5.15) \quad f_\varepsilon(I, \theta) = \left((I - \omega_*)e^{-2\pi\lambda} + \omega_*, \theta + \frac{1}{\lambda}(I - \omega_*)(1 - e^{-2\pi\lambda}) + 2\pi\omega_* \right),$$

which is the first-return map to the section $\Lambda_\varepsilon = \tilde{\Lambda}_\varepsilon \cap \{s = 0 \pmod{2\pi}\}$.

In particular, for $I = \omega_*$ we have $f_\varepsilon(I, \theta) = (\omega_*, \theta + 2\pi\omega_*) = (I, \theta + 2\pi\omega_*)$. That is, $A_\varepsilon = \tilde{A}_\varepsilon \cap \{s = 0 \pmod{2\pi}\}$ is an invariant circle for f_ε of irrational rotation number $2\pi\omega_*$.

From (5.14) we have that, for (I, θ) with $\theta = \theta_0 - \frac{1}{\lambda}(I - \omega_*)$,

$$(5.16) \quad \|f_\varepsilon^k(I, \theta) - f_\varepsilon^k(\omega_*, \theta_0)\| \leq \left(1 + \frac{1}{\lambda^2}\right)^{1/2} |I - \omega_*| e^{-2\pi\lambda k} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

This shows that A_ε is a global attractor for the map f_ε on Λ_ε , and, moreover, the orbits of (I, θ) and (ω_*, θ_0) become asymptotically close to one another as $n \rightarrow \infty$.

We have

$$Df_\varepsilon(I, \theta) = \begin{pmatrix} e^{-2\pi\lambda} & 0 \\ \frac{1}{\lambda}(1 - e^{-2\pi\lambda}) & 1 \end{pmatrix}$$

with eigenvalues $e^{-2\pi\lambda}$ and 1 and with corresponding eigenvectors

$$\begin{pmatrix} -\lambda \\ 1 \end{pmatrix} \text{ and } \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

respectively. The eigenvalue 1 is associated to the dynamics along the θ -coordinate, and the eigenvalue of $e^{-2\pi\lambda} < 1$ is associated to the dynamics along the I -coordinate.

We conclude that A_ε is a NHIM for $(f_\varepsilon)|_{\Lambda_\varepsilon}$, for which there is only stable manifold $W^s(A_\varepsilon)$ tangent to $\begin{pmatrix} -\lambda \\ 1 \end{pmatrix}$, and no unstable manifold $W^u(A_\varepsilon)$. We note that for $\lambda \gtrsim 0$ (recall that $\lambda = \varepsilon\rho$) the Lyapunov multipliers 1 and $e^{-2\pi\lambda} \lesssim 1$ for $(f_\varepsilon)|_{\Lambda_\varepsilon}$, are dominated by the contraction rate of Df_ε on the stable bundle E^s of Λ_ε , which is $e^{-2\pi+O(\lambda)}$; see Section 5.3.

Since $|\det(Df_\varepsilon)| = e^{-2\pi\lambda} < 1$ we have that f_ε is area-contracting on Λ_ε , hence it is conformally symplectic, i.e.

$$(5.17) \quad (f_\varepsilon)^*_{|\Lambda_\varepsilon}(\omega|_{\Lambda_\varepsilon}) = e^{-2\pi\lambda}\omega|_{\Lambda_\varepsilon}.$$

We now show that f_ε is a λ -perturbation of f_0 , a time- 2π map for the rotator part of the unperturbed system given in (5.6). Since

$$e^{-2\pi\lambda} = 1 - 2\pi\lambda + O(\lambda^2),$$

we have

$$f_\varepsilon(I, \theta) = (I - 2\pi\lambda(I - \omega_*) + O(\lambda^2), \theta + 2\pi I - 2\pi^2\lambda(I - \omega_*) + O(\lambda^2)).$$

Therefore f_ε is a λ -perturbation of f_0 in (5.6), i.e.

$$f_\varepsilon(I, \theta) = f_0(I, \theta) + O(\lambda) = f_0(I, \theta) + O(\varepsilon\rho).$$

5.8. The case of non-vanishing perturbation. In this case the time-periodic perturbation of the Hamiltonian in (3.5), is of the form

$$(5.18) \quad H_1(p, q, I, \theta, s) = \cos q \cdot g(\theta, s).$$

The perturbation H_1 does not vanish at the hyperbolic fixed point of the pendulum $(p, q) = (0, 0)$. The dissipative perturbation is given by the vector field $\varepsilon \mathcal{X}_\rho$, where \mathcal{X}_ρ is given by (3.7), as before (see (5.9) and (5.10)).

From (3.9), since $f'(q) = -\sin q$ vanishes at $q = 0$, we obtain that the unperturbed NHIM $\tilde{\Lambda}_0$ survives the perturbation, that is $\tilde{\Lambda}_\varepsilon = \tilde{\Lambda}_0$ for all ε .

When $\lambda \neq 0$, the perturbed dynamics restricted to $\tilde{\Lambda}_\varepsilon = \tilde{\Lambda}_0$ is given by the following equations:

$$(5.19) \quad \begin{cases} \dot{I} = -\lambda(I - \omega_*) - \varepsilon \frac{\partial g}{\partial \theta}(\theta, s) \\ \dot{\theta} = I \\ \dot{s} = 1. \end{cases}$$

Using the expression of g in (3.11), this system can be reduced to the second-order nonlinear differential equation

$$\ddot{\theta} + \lambda \dot{\theta} - \varepsilon a_{10} \sin \theta - \lambda \omega_* = 0.$$

Ignoring the last term, the remaining terms represent the equation of the damped non-linear pendulum, for which explicit solutions are unknown; an analytical approximation can be found in [Joh14]. Hence, we do not have an explicit formula for the time- 2π map f_ε in this case.

6. Existence of a Transverse Homoclinic Intersection. In the sequel, we will identify vector fields with differential operators, which is a standard operation in differential geometry (see, e.g., [BG05]). That is, given a smooth vector field \mathcal{X} and a smooth function f on the manifold M , we denote:

$$(6.1) \quad (\mathcal{X}f)(z) = \sum_j (\mathcal{X})_j(z) \frac{\partial f}{\partial z_j}(z),$$

where $z_j, j \in \{1, \dots, \dim(M)\}$, are local coordinates. Similarly, a smooth time-dependent and parameter-dependent vector field acts as a differential operator by

$$(6.2) \quad (\mathcal{X}f)(z, t; \varepsilon) = \sum_j (\mathcal{X})_j(z, t; \varepsilon) \frac{\partial f}{\partial z_j}(z).$$

For the pendulum system, whose hamiltonian h_1 is given in (3.3), we denote by $(p_0(t), q_0(t))$ a parametrization of a separatrix of the pendulum, with $(p_0(0), q_0(0)) = (p_0, q_0)$, where (p_0, q_0) is some initial point; this parametrization is explicitly given in (3.4). We define a new locally defined system of symplectic coordinates (y, x) in a neighborhood of the separatrix – chosen away from the hyperbolic equilibrium point – as follows. The coordinate y is chosen to be equal to the energy of the pendulum, i.e.,

$$(6.3) \quad y = h_1(p, q) = \frac{p^2}{2} + (\cos(q) - 1)$$

and is defined in a whole neighborhood of one of its separatrices. The coordinate x is defined by

$$dx = \frac{dt}{\|\nabla y\|},$$

where $dt = (dp^2 + dq^2)^{\frac{1}{2}}$. It is immediate to see that x equals the time τ it takes the solution $(p(t), q(t))$ to go along the y -level set from one point to another (see [GdlLM21]). This coordinate system (y, x) constructed above is not defined in a neighborhood of the separatrix that contains the hyperbolic equilibrium point, since this is a critical point of the energy function. We define this coordinate system only in some neighborhood \mathcal{N} of a segment of the separatrix containing (p_0, q_0) . On this neighborhood, we have $dy \wedge dx = dp \wedge dq$. Relative to this new coordinate system, the separatrix is given by $y = 0$.

An arbitrary point on the separatrix can be given in terms of the (p, q) -coordinates as $(p_0(\tau), q_0(\tau))$ for some $\tau \in \mathbb{R}$, and in terms of the (y, x) -coordinates as $(0, x)$ for some $x \in \mathbb{R}$, where $x = \tau$.

Now let's extend this coordinate system to a system of coordinates (y, x, I, θ, s) on some neighborhood $\tilde{\mathcal{N}}$ of $\{(p_0(\tau), q_0(\tau), I, \theta, s)\}$ in the extended phase space.

Relative to this coordinate system, in the unperturbed case, the stable/unstable manifolds $W^s(\tilde{\Lambda}_0) = W^u(\tilde{\Lambda}_0)$ are locally given by $y = 0$. A point $\tilde{z}_0 \in W^s(\tilde{\Lambda}_0) = W^u(\tilde{\Lambda}_0)$ can be written in terms of the original coordinates (p, q, I, θ, s) as

$$\tilde{z}_0 = (p_0(\tau), q_0(\tau), I, \theta, s), \text{ for some } \tau \in \mathbb{R},$$

and in terms of the extended coordinates (y, x, I, θ, s) as

$$\tilde{z}_0 = (0, x, I, \theta, s), \text{ for } x = \tau \in \mathbb{R}.$$

When we apply the flow to the point \tilde{z}_0 we obtain

$$\tilde{\Phi}_0^t(\tilde{z}_0) = (p_0(\tau + t), q_0(\tau + t), I, \theta + \omega(I)t, s + t).$$

Observe that if we denote by $\tilde{z}_0^\pm := (p, q, I, \theta, s) = (0, 0, I, \theta, s)$, we have $\tilde{\Phi}_0^t(\tilde{z}_0^\pm) = (0, 0, I, \theta + \omega(I)t, s + t)$, therefore:

$$\tilde{\Phi}_0^t(\tilde{z}_0) - \tilde{\Phi}_0^t(\tilde{z}_0^\pm) \rightarrow 0, \quad \text{as } t \rightarrow \pm\infty.$$

In the perturbed case, for $\varepsilon \neq 0$ small and $\lambda = \rho\varepsilon$, we can locally describe both the stable and unstable manifolds of $\tilde{\Lambda}_\varepsilon$ as graphs of $C^{\ell-1}$ -smooth functions $y_\varepsilon^s, y_\varepsilon^u$, over (x, I, θ, s) , recalling that $x = \tau$, given by

$$\begin{aligned} y_\varepsilon^s &= y_\varepsilon^s(x, I, \theta, s; \rho) = y_\varepsilon^s(\tau, I, \theta, s; \rho), \\ y_\varepsilon^u &= y_\varepsilon^u(x, I, \theta, s; \rho) = y_\varepsilon^u(\tau, I, \theta, s; \rho), \end{aligned}$$

respectively, for $(0, x, I, \theta, s) \in \tilde{\mathcal{N}}$. We stress the dependence of ρ of these functions because will be important in the sequel.

Observe that, when $\varepsilon = 0$ we have the equation of the separatrix of the pendulum

$$y_0^s(\tau, I, \theta, s; \rho) = y_0^u(\tau, I, \theta, s; \rho) = 0.$$

Consequently $y_\varepsilon^u, y_\varepsilon^s = O(\varepsilon)$.

We recall the following Melnikov-type result for non-conservative perturbations:

Theorem 6.1 (Splitting of the Stable and Unstable Manifolds [GdlLM21]).

Fix $\rho_0 > 0$, then there exists $\varepsilon_0 > 0$ such that for any $0 \leq \rho \leq \rho_0$ and $0 \leq |\varepsilon| \leq \varepsilon_0$ we have: for $(0, \tau, I, \theta, s) \in \tilde{\mathcal{N}}$, the difference between $y_\varepsilon^s(\tau, I, \theta, s; \rho)$ and $y_\varepsilon^u(\tau, I, \theta, s; \rho)$ is given by

$$y_\varepsilon^s - y_\varepsilon^u = -\varepsilon \int_{-\infty}^{+\infty} ((\mathcal{X}^1 h_1)(\tilde{\Phi}_0^t(\tilde{z}_0)) - (\mathcal{X}^1 h_1)(\tilde{\Phi}_0^t(\tilde{z}_0^\pm))) dt + O(\varepsilon^2),$$

where we recall that $h_1(p, q) = \frac{p^2}{2} + (\cos q - 1)$ and \mathcal{X}^1 is given in (5.10).

Corollary 6.2 (Sufficient Conditions for the Existence of a Transverse Homoclinic Intersection).

Fix $\rho_0 > 0$, then there exists $\varepsilon_0 > 0$ such that for any $0 \leq \rho \leq \rho_0$ and $|\varepsilon| \leq \varepsilon_0$ we have: for $(0, \tau, I, \theta, s) \in \tilde{\mathcal{N}}$, the difference between $y_\varepsilon^s(\tau, I, \theta, s; \rho)$ and $y_\varepsilon^u(\tau, I, \theta, s; \rho)$ is given by

$$(6.4) \quad y_\varepsilon^s - y_\varepsilon^u = -\varepsilon \left[\int_{-\infty}^{+\infty} \{h_1, H_1\}(p_0(\tau+t), q_0(\tau+t), I, \theta + \omega(I)t, s+t) dt \right. \\ \left. - \rho \int_{-\infty}^{+\infty} p_0^2(t) dt \right] + O(\varepsilon^2),$$

where $\{\cdot, \cdot\}$ denotes the Poisson bracket.

If $\tau^* = \tau^*(I, \theta, s)$ is a non-degenerate zero of the mapping

$$(6.5) \quad \tau \in \mathbb{R} \mapsto - \int_{-\infty}^{+\infty} \{h_1, H_1\}(p_0(\tau+t), q_0(\tau+t), I, \theta + \omega(I)t, s+t) dt$$

then there exists $0 < \rho_1 \leq \rho_0$ such that for all $0 \leq \rho \leq \rho_1$

$$(6.6) \quad \tau \in \mathbb{R} \mapsto - \left[\int_{-\infty}^{+\infty} \{h_1, H_1\}(p_0(\tau+t), q_0(\tau+t), I, \theta + \omega(I)t, s+t) dt \right. \\ \left. - \rho \int_{-\infty}^{+\infty} p_0^2(t) dt \right]$$

has a non degenerate zero $\tau^*(I, \theta, s; \rho)$.

Moreover, there exists $0 < \varepsilon_1 \leq \varepsilon_0$ such that for all $0 \leq \rho \leq \rho_1$ and $0 < |\varepsilon| \leq \varepsilon_1$, $W^s(\tilde{\Lambda}_\varepsilon)$ and $W^u(\tilde{\Lambda}_\varepsilon)$ have a transverse homoclinic intersection which can be parametrized as

$$(6.7) \quad (\tau^*, y_\varepsilon^s(\tau^*, I, \theta, s; \rho), I, \theta, s) = (\tau^*, y_\varepsilon^u(\tau^*, I, \theta, s; \rho), I, \theta, s),$$

where $\tau^* = \tau^*(I, \theta, s; \rho, \varepsilon) = \tau^*(I, \theta, s; \rho) + O(\varepsilon) = \tau^*(I, \theta, s) + O(\rho, \varepsilon)$, for (I, θ, s) in some open set in $\tilde{U} \subseteq \mathbb{R} \times \mathbb{T}^1 \times \mathbb{T}^1$.

Proof. From Theorem 6.1, we have

$$y_\varepsilon^s - y_\varepsilon^u = -\varepsilon \int_{-\infty}^{+\infty} ((\mathcal{X}^1 h_1)(\tilde{\Phi}_0^t(\tilde{z}_0)) - (\mathcal{X}^1 h_1)(\tilde{\Phi}_0^t(\tilde{z}_0^\pm))) dt + O(\varepsilon^2).$$

446 As $\lambda = \varepsilon\rho$, the vector field \mathcal{X}^1 (see (5.10)) is the sum of the Hamiltonian vector field $J\nabla H_1$
 447 and the dissipative vector field

$$448 \quad \mathcal{X}_\rho(p, q, I, \theta) = (-\rho p, 0, -\rho(I - \omega_*), 0),$$

therefore

$$\begin{aligned} y_\varepsilon^s - y_\varepsilon^u &= -\varepsilon \int_{-\infty}^{+\infty} \left[(J\nabla H_1 + \mathcal{X}_\rho)h_1(\tilde{\Phi}_0^t(\tilde{z}_0)) - (J\nabla H_1 + \mathcal{X}_\rho)h_1(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right] dt + O(\varepsilon^2) \\ &= -\varepsilon \left[\int_{-\infty}^{+\infty} (J\nabla H_1 h_1)(\tilde{\Phi}_0^t(\tilde{z}_0)) - (J\nabla H_1 h_1)(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) dt \right. \\ &\quad \left. + \int_{-\infty}^{+\infty} (\mathcal{X}_\rho h_1)(\tilde{\Phi}_0^t(\tilde{z}_0)) - (\mathcal{X}_\rho h_1)(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) dt \right] + O(\varepsilon^2) \\ &:= -\varepsilon(\mathcal{F}_1 + \mathcal{F}_2) + O(\varepsilon^2). \end{aligned}$$

In the above,

$$\begin{aligned} \mathcal{F}_1 &:= \int_{-\infty}^{+\infty} (J\nabla H_1 h_1)(\tilde{\Phi}_0^t(\tilde{z}_0)) - (J\nabla H_1 h_1)(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) dt \\ &= \int_{-\infty}^{+\infty} \{h_1, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0)) - \{h_1, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) dt, \\ &= \int_{-\infty}^{+\infty} \left(\{h_1, H_1\}(p_0(\tau + t), q_0(\tau + t), I, \theta + \omega(I)t, s + t) \right. \\ &\quad \left. - \{h_1, H_1\}(0, 0, I, \theta + \omega(I), s + t) \right) dt \\ \mathcal{F}_2 &:= \int_{-\infty}^{+\infty} (\mathcal{X}_\rho h_1)(\tilde{\Phi}_0^t(\tilde{z}_0)) - (\mathcal{X}_\rho h_1)(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) dt \\ &= \rho \int_{-\infty}^{+\infty} (\mathcal{X}_{\omega_*} h_1)(\tilde{\Phi}_0^t(\tilde{z}_0)) - (\mathcal{X}_{\omega_*} h_1)(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) dt \end{aligned}$$

where we denote $\mathcal{X}_{\omega_*} = (-p, 0, -(I - \omega_*), 0)$. Since $\mathcal{X}_{\omega_*} h_1 = -p \frac{\partial h_1}{\partial p} = -p^2$, and recalling that $\tilde{\Phi}_0^t(\tilde{z}_0) = (p_0(\tau + t), q_0(\tau + t), I, \theta + \omega(I)t, s + t)$ and $\tilde{\Phi}_0^t(\tilde{z}_0^\pm) = (0, 0, I, \theta + \omega(I)t, s + t)$, we obtain

$$\mathcal{F}_2 = -\rho \int_{-\infty}^{+\infty} p_0^2(\tilde{\Phi}_0^t(\tilde{z}_0)) dt = -\rho \int_{-\infty}^{+\infty} p_0^2(\tau + t) dt$$

Finally,

$$\begin{aligned} y_\varepsilon^s - y_\varepsilon^u &= -\varepsilon \int_{-\infty}^{+\infty} \left(\{h_1, H_1\}(p_0(\tau+t), q_0(\tau+t), I, \theta + \omega(I)t, s+t) \right. \\ &\quad \left. - \{h_1, H_1\}(0, 0, I, \theta + \omega(I), s+t) \right) dt \\ &\quad + \varepsilon \rho \int_{-\infty}^{+\infty} p_0^2(\tau+t) dt + O(\varepsilon^2). \end{aligned}$$

449 Note that $\{h_1, H_1\} = -\sin q \frac{\partial H_1}{\partial p} - p \frac{\partial H_1}{\partial q}$ hence $\{h_1, H_1\}(0, 0, I, \theta + \omega(I), s+t) = 0$. Also note
 450 that by the change of variable formula $\int_{-\infty}^{+\infty} p_0^2(\tau+t) dt = \int_{-\infty}^{+\infty} p_0^2(t) dt$. Thus, we obtain the
 451 first part of Corollary 6.2.

The second part of Corollary 6.2 is as follows. First, if $\tau^* = \tau^*(I, \theta, s)$ is a non-degenerate zero of the mapping (6.5), there exists $0 < \rho_1 \leq \rho_0$ such that the function:

$$\tau \in \mathbb{R} \mapsto - \left[\int_{-\infty}^{+\infty} \{h_1, H_1\}(p_0(\tau+t), q_0(\tau+t), I, \theta + \omega(I)t, s+t) dt - \rho \int_{-\infty}^{+\infty} p_0^2(t) dt \right]$$

452 also has a non-degenerate zero $\tau^*(I, \theta, s; \rho) = \tau^*(I, \theta, s) + O(\rho)$ for any $0 \leq \rho \leq \rho_1$.

453 Now, we apply the implicit function theorem to find the zeroes of the function:

$$454 \quad \tau \rightarrow y_\varepsilon^s(\tau, I, \theta, s; \rho) - y_\varepsilon^u(\tau, I, \theta, s; \rho),$$

455 obtaining a value $0 < \tilde{\varepsilon}_1(\rho) \leq \varepsilon_0$, such that, for any $0 < \varepsilon \leq \tilde{\varepsilon}_1$, this map has a non-degenerate
 456 zero $\tau^*(I, \theta, s; \rho, \varepsilon) = \tau^*(I, \theta, s; \rho) + O(\varepsilon) = \tau^*(I, \theta, s) + O(\rho, \varepsilon)$. An important observation is
 457 that $\tilde{\varepsilon}_1(0) \neq 0$, therefore we set $\varepsilon_1 = \min_{[0, \rho_1]} \tilde{\varepsilon}_1(\rho) > 0$. In this way, arguing as in [DLS06], the
 458 stable and unstable manifolds $W^s(\tilde{\Lambda}_\varepsilon)$ and $W^u(\tilde{\Lambda}_\varepsilon)$ have a transverse homoclinic intersection
 459 which can be parametrized as in (6.7). ■

460 Provided that the unperturbed stable and unstable manifolds of the NHIM coincide,
 461 adding a generic Hamiltonian perturbation makes the stable and unstable manifolds to inter-
 462 sect transversally; see, e.g., [GdlL18]. However, non-conservative perturbations can in general
 463 destroy the homoclinic intersection; this is for example the case of the dissipative pendulum
 464 shown in Fig. 2b. In contrast, the Corollary 6.2 shows that for the system (3.8), where the
 465 dissipation is of the same order as the forcing, that is $\lambda = \varepsilon \rho$, the perturbed stable and unsta-
 466 ble manifolds intersect transversally for all sufficiently small perturbation parameter values ρ .
 467 Later, in Section 8, we will be interested in taking $\rho = \rho(\varepsilon) = \frac{\bar{\rho}}{\log(\frac{1}{\varepsilon})}$, but, clearly, for ε small
 468 enough, these values of ρ satisfy the hypotheses of Corollary 6.2. The result is summarized in
 469 next corollary.

470 **Corollary 6.3 (Existence of Transverse Intersection in the Model).** *Take any $\bar{\rho} > 0$. Consider*
 471 *the perturbation H_1 given by (3.10) or (3.11) and the dissipative perturbation as in (3.7) with*
 472 *$\lambda = \varepsilon \frac{\bar{\rho}}{\log(\frac{1}{\varepsilon})}$. Then there exists ε_0 sufficiently small such that for all $0 < |\varepsilon| < \varepsilon_0$, $W^s(\tilde{\Lambda}_\varepsilon)$ and*
 473 *$W^u(\tilde{\Lambda}_\varepsilon)$ have a transverse homoclinic intersection $\tilde{\Gamma}_\varepsilon$ which can be parametrized as in (6.7).*

474 **Proof.** The proof follows by the fact that in this case $\rho = \frac{\bar{\rho}}{\log(\frac{1}{\varepsilon})}$ satisfies the conditions of
 475 Corollary 6.2 if ε is small enough. ■

7. Computation of the scattering map for the perturbed system.

7.1. The scattering map. We give a brief description of the scattering map, following [DdlLS08]. Consider the general case of a normally hyperbolic invariant manifold Λ for a flow Φ^t on some smooth manifold M . Let $W^s(\Lambda)$, $W^u(\Lambda)$ be the stable and unstable manifolds of Λ . First, let

$$\begin{aligned}\Omega^+ : W^s(\Lambda) &\rightarrow \Lambda, \\ \Omega^- : W^u(\Lambda) &\rightarrow \Lambda,\end{aligned}$$

be the canonical projections along fibers, assigning to each point $x \in W^s(\Lambda)$ its stable foot point $x^+ = \Omega^+(x)$, uniquely defined by $x \in W^s(x^+)$, and, similarly, assigning to $x \in W^u(\Lambda)$ its unstable footpoint $x^- = \Omega^-(x)$ uniquely defined by $x \in W^u(x^-)$.

Second, choose and fix a ‘homoclinic channel’, which is a homoclinic manifold Γ in $W^u(\Lambda) \cap W^s(\Lambda)$ that satisfies the following strong transversality conditions:

$$\begin{aligned}T_x \Gamma &= T_x W^s(\Lambda) \cap T_x W^u(\Lambda), \\ T_x M &= T_x \Gamma \oplus T_x W^u(x^-) \oplus T_x W^s(x^+),\end{aligned}$$

for all $x \in \Gamma$, and such that

$$\Omega_{|\Gamma}^\pm : \Gamma \rightarrow \Omega^\pm(\Gamma) \text{ is a diffeomorphism.}$$

Then, the scattering map associated to the homoclinic channel Γ is the mapping $\sigma : \Omega^-(\Gamma) \rightarrow \Omega^+(\Gamma)$ defined by

$$\sigma = \Omega^+ \circ (\Omega^-)^{-1}.$$

The map σ is a locally defined diffeomorphism on Λ . Moreover, σ is symplectic provided that M , Λ , Φ^t are symplectic.

Remark 7.1. We have $\sigma(x^-) = x^+$ if and only if

$$(7.1) \quad d(\Phi^{-T_-}(x), \Phi^{-T_-}(x^-)) \rightarrow 0, \text{ and } d(\Phi^{T_+}(x), \Phi^{T_+}(x^+)) \rightarrow 0$$

as $T_-, T_+ \rightarrow +\infty$, respectively, for some uniquely defined $x \in \Gamma$. This means that for orbits in Λ of the form $x'_{\text{end}} = \Phi^{T_+} \circ \sigma \circ \Phi^{T_-}(x'_{\text{start}})$, where $x'_{\text{start}} = \Phi^{-T_-}(x^-)$ and $x'_{\text{end}} = \Phi^{T_+}(x^+)$, one can find homoclinic orbit segments in M of the form $x_{\text{end}} = \Phi^{T_+ + T_-}(x_{\text{start}})$, such that x_{start} is arbitrarily close to x'_{start} and x_{end} is arbitrarily close to x'_{end} . See Fig. 4.

7.2. The scattering map of the perturbed system. Assuming that the conditions in Corollary 6.2 are satisfied, then $W^s(\tilde{\Lambda}_\varepsilon)$ and $W^u(\tilde{\Lambda}_\varepsilon)$ intersect transversally in the homoclinic channel $\tilde{\Gamma}_\varepsilon$, which can be parametrized as in Corollary 6.2, for all $0 < |\varepsilon| < \varepsilon_1$. Let $\tilde{z}_\varepsilon \in \tilde{\Gamma}_\varepsilon$ be a homoclinic point for the perturbed extended flow $\tilde{\Phi}_\varepsilon^t$. In terms of the coordinates from Section 6, we have

$$\tilde{z}_\varepsilon = (\tau^*(I, \theta, s; \rho, \varepsilon), y_\varepsilon^s(\tau^*(I, \theta, s; \rho, \varepsilon), I, \theta, s; \rho), I, \theta, s)$$

where $\tau^*(I, \theta, s; \rho, \varepsilon) = \tau^*(I, \theta, s; \rho) + O(\varepsilon)$ and $\tau^*(I, \theta, s; \rho) = \tau^*(I, \theta, s) + O(\varepsilon)$ is a non-degenerate zero of the mapping (6.6) near $\tau^*(I, \theta, s)$, a chosen non-degenerate zero of the mapping (6.5).




Figure 4. Homoclinic orbit segment approximating an orbit obtained by applying the scattering map and the inner map.

Because of the smooth dependence of the NHIM and of its stable and unstable manifolds on the perturbation parameter, to the homoclinic point $\tilde{z}_\varepsilon = (z_\varepsilon, s)$ for perturbed flow $\tilde{\Phi}_\varepsilon^t$ it corresponds a homoclinic point $\tilde{z}_0 = (z_0, s)$ for the unperturbed flow $\tilde{\Phi}_0^t$, which is $O(\varepsilon)$ -close to \tilde{z}_ε . In fact, going back to the original coordinates, the point \tilde{z}_ε becomes:

$$\begin{aligned}
 \tilde{z}_\varepsilon &= (p_0(\tau^*(I, \theta, s; \rho, \varepsilon)), q_0(\tau^*(I, \theta, s; \rho, \varepsilon)), I, \theta, s) + O(\varepsilon) \\
 &= (p_0(\tau^*(I, \theta, s; \rho)), q_0(\tau^*(I, \theta, s; \rho)), I, \theta, s) + O(\varepsilon) \\
 &= \tilde{z}_0 + O(\varepsilon), \text{ where} \\
 \tilde{z}_0 &= (p_0(\tau^*(I, \theta, s; \rho)), q_0(\tau^*(I, \theta, s; \rho)), I, \theta, s)
 \end{aligned}
 \tag{7.2}$$

Note that in the above the $O(\varepsilon)$ -error only affects the p, q components.

We denote the stable- and unstable-footpoints of \tilde{z}_ε and of \tilde{z}_0 by $\tilde{z}_\varepsilon^\pm$ and \tilde{z}_0^\pm , respectively. Recall that we already know that $\tilde{z}_0^\pm = (0, 0, I, \theta, s)$. Summarizing the notation:

- $\tilde{z}_\varepsilon \in \tilde{\Gamma}_\varepsilon \subset W^s(\tilde{\Lambda}_\varepsilon) \pitchfork W^u(\tilde{\Lambda}_\varepsilon)$;
- $\tilde{z}_\varepsilon^\pm = \Omega^\pm(\tilde{z}_\varepsilon) \in \tilde{\Lambda}_\varepsilon$;
- $\tilde{z}_0 \in \tilde{\Gamma}_0 \subset W^s(\tilde{\Lambda}_0) \pitchfork W^u(\tilde{\Lambda}_0)$;
- $\tilde{z}_0^\pm = \Omega^\pm(\tilde{z}_0) \in \tilde{\Lambda}_0$;

Under the above assumptions, we have $\tilde{\sigma}_\varepsilon(\tilde{z}_\varepsilon^-) = \tilde{z}_\varepsilon^+$, and $\tilde{\sigma}_0(\tilde{z}_0^-) = \tilde{z}_0^+$. We recall that, in our model, for the unperturbed system, $\tilde{z}_0^- = \tilde{z}_0^+ = \tilde{z}_0^\pm$ and therefore the scattering map is the identity: $\tilde{\sigma}_0 = \text{Id}$.

The perturbed scattering map $\tilde{\sigma}_\varepsilon$ can be expanded in terms of powers of ε , with the zero-th order term being the unperturbed scattering map $\tilde{\sigma}_0$, as follows

$$\begin{aligned}
 \tilde{\sigma}_\varepsilon(I, \theta, s) &= \tilde{\sigma}_0(I, \theta, s) + \varepsilon \mathcal{S}(I, \theta, s) + O(\varepsilon^2) \\
 &= (I, \theta, s) + \varepsilon \mathcal{S}(I, \theta, s) + O(\varepsilon^2),
 \end{aligned}$$

where $\mathcal{S} = (\mathcal{S}^I, \mathcal{S}^\theta, \text{Id}^s)$.

In the sequel, we evaluate the components \mathcal{S}^I and \mathcal{S}^θ in order to compute the change in action I and the change in angle θ by the scattering map. We follow the approach in [GdlLM21, GdlLM22].

7.3. Change in Action by the Scattering Map. We use the following result:

Theorem 7.2 (Change in Action by the Scattering Map [GdlLM21]). *For a general non-conservative perturbation \mathcal{X}^1 of (3.1) like in (5.9), the change in action I by the scattering map $\tilde{\sigma}_\varepsilon$ is given by:*

$$(7.3) \quad I(\tilde{z}_\varepsilon^+) - I(\tilde{z}_\varepsilon^-) = \varepsilon \int_{-\infty}^{+\infty} \left(\mathcal{X}^1 I(\tilde{\Phi}_0^t(\tilde{z}_0)) - \mathcal{X}^1 I(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt + O(\varepsilon^2),$$

where \tilde{z}_ε and \tilde{z}_0 are given in (7.2), and we denote $I_0 = I(\tilde{z}_0) = I(\tilde{z}_0^\pm)$.

Denote $I(\tilde{z}_\varepsilon^\pm) = I_\varepsilon^\pm$. Applying Theorem 7.2 in the case of (5.10):

$$\begin{aligned} \mathcal{X}^1 &= J\nabla H_1 + (-\rho p, 0, -\rho(I - \omega_*), 0) \\ &= J\nabla H_1 + \mathcal{X}_\rho \end{aligned}$$

we obtain

$$\begin{aligned} I_\varepsilon^+ - I_\varepsilon^- &= \varepsilon \int_{-\infty}^{\infty} \left((J\nabla H_1 + \mathcal{X}_\rho) I(\tilde{\Phi}_0^t(\tilde{z}_0)) - (J\nabla H_1 + \mathcal{X}_\rho) I(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt + O(\varepsilon^2) \\ &= \varepsilon \int_{-\infty}^{\infty} \left((J\nabla H_1 I)(\tilde{\Phi}_0^t(\tilde{z}_0)) - (J\nabla H_1 I)(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt \\ &\quad + \varepsilon \int_{-\infty}^{\infty} \left(\mathcal{X}_\rho I(\tilde{\Phi}_0^t(\tilde{z}_0)) - \mathcal{X}_\rho I(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt + O(\varepsilon^2) \\ &= \varepsilon(\mathcal{S}_1^I + \mathcal{S}_2^I) + O(\varepsilon^2) \end{aligned}$$

where

$$\begin{aligned} \mathcal{S}_1^I &:= \int_{-\infty}^{\infty} \left(\{I, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0)) - \{I, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt \\ &= \int_{-\infty}^{+\infty} \left(\{I, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \right. \\ &\quad \left. - \{I, H_1\}(0, 0, I, \theta + \omega(I)t, s + t) \right) dt \\ \mathcal{S}_2^I &= \int_{-\infty}^{\infty} \left(\mathcal{X}_\rho I(\tilde{\Phi}_0^t(\tilde{z}_0)) - \mathcal{X}_\rho I(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt, \end{aligned}$$

where $\tau^* = \tau^*(I, \theta, s; \rho)$ is a non-degenerate zero of the function (6.6).

Since $\mathcal{X}_\rho I = -\rho(I - \omega_*)$, $\tilde{\Phi}_0^t(\tilde{z}_0) = (p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t)$ and $\tilde{\Phi}_0^t(\tilde{z}_0^\pm) = (0, 0, I, \theta + \omega(I)t, s + t)$, we have

$$\mathcal{S}_2^I := \int_{-\infty}^{\infty} \left(-\rho I(\tilde{\Phi}_0^t(\tilde{z}_0)) + \rho I(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt = 0.$$

Thus, we have proved the following result:

Corollary 7.3. For the perturbation $\mathcal{X}^1 = J\nabla H_1 + \mathcal{X}_\rho$,

$$(7.4) \quad I_\varepsilon^+ - I_\varepsilon^- = \varepsilon \int_{-\infty}^{+\infty} (\{I, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \\ - \{I, H_1\}(0, 0, I, \theta + \omega(I)t, s + t)) dt + O(\varepsilon^2)$$

where $\tau^* = \tau^*(I, \theta, s; \rho)$ is a non-degenerate zero of the function (6.6).

In the case when H_1 is as in (3.10) or (3.11), $\{I, H_1\} = -\frac{\partial H_1}{\partial \theta} = a_{10}f(q)\sin\theta$, where $f(q) = \cos q - 1$ or $f(q) = \cos q$, so

$$(7.5) \quad I_\varepsilon^+ - I_\varepsilon^- = \varepsilon a_{10} \int_{-\infty}^{+\infty} (\cos(q_0(\tau^* + t)) - 1) \sin(\theta + \omega(I)t) dt + O(\varepsilon^2).$$

7.4. Change in Angle by the Scattering Map. We use the following result:

Theorem 7.4 (Change in Angle by the Scattering Map [GdlLM21]). For a general non-conservative perturbation \mathcal{X}^1 of (3.1) like in (5.9), the change in angle θ by the scattering map $\tilde{\sigma}_\varepsilon$ is given by:

$$(7.6) \quad \theta(\tilde{z}_\varepsilon^+) - \theta(\tilde{z}_\varepsilon^-) = \varepsilon \int_{-\infty}^{+\infty} \mathcal{X}^1 \theta(\tilde{\Phi}_0^t(\tilde{z}_0)) - \mathcal{X}^1 \theta(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) dt \\ - \varepsilon \int_{-\infty}^{+\infty} (\mathcal{X}^1 I(\tilde{\Phi}_0^t(\tilde{z}_0)) - \mathcal{X}^1 I(\tilde{\Phi}_0^t(\tilde{z}_0^\pm))) t dt \cdot \left(\frac{\partial^2 h_0}{\partial I^2}(I_0) \right) \\ + O(\varepsilon^2),$$

where \tilde{z}_ε and \tilde{z}_0 are given in (7.2), and we denote $I_0 = I(\tilde{z}_0) = I(\tilde{z}_0^\pm)$.

Denote $\theta(\tilde{z}_\varepsilon^\pm) = \theta_\varepsilon^\pm$. Applying Theorem 7.4 in the case of (5.10), i.e., $\mathcal{X}^1 = J\nabla H_1 + \mathcal{X}_\rho$ we obtain

$$(7.7) \quad \theta_\varepsilon^+ - \theta_\varepsilon^- = \varepsilon \int_{-\infty}^{+\infty} \left((J\nabla H_1 + \mathcal{X}_\rho) \theta(\tilde{\Phi}_0^t(\tilde{z}_0)) - (J\nabla H_1 + \mathcal{X}_\rho) \theta(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt \\ - \varepsilon \int_{-\infty}^{+\infty} \left((J\nabla H_1 + \mathcal{X}_\rho) I(\tilde{\Phi}_0^t(\tilde{z}_0)) \right. \\ \left. - (J\nabla H_1 + \mathcal{X}_\rho) I(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) t dt \cdot \left(\frac{\partial^2 h_0}{\partial I^2}(I) \right) \\ + O(\varepsilon^2)$$

We simplify the first integral above by splitting into two integrals:

$$\varepsilon \left[\int_{-\infty}^{+\infty} \left((J\nabla H_1 \theta)(\tilde{\Phi}_0^t(\tilde{z}_0)) - (J\nabla H_1 \theta)(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt \right. \\ \left. + \int_{-\infty}^{+\infty} \left(\mathcal{X}_\rho \theta(\tilde{\Phi}_0^t(\tilde{z}_0)) - \mathcal{X}_\rho \theta(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt \right] \\ = \varepsilon (\mathcal{S}_1^\theta + \mathcal{S}_2^\theta),$$

where

$$\begin{aligned}\mathcal{S}_1^\theta &:= \int_{-\infty}^{\infty} \left(\{\theta, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0)) - \{\theta, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt \\ &= \int_{-\infty}^{\infty} \left(\{\theta, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \right. \\ &\quad \left. - \{\theta, H_1\}(0, 0, I, \theta + \omega(I)t, s + t) \right) dt, \\ \mathcal{S}_2^\theta &= \int_{-\infty}^{\infty} \left(\mathcal{X}_\rho \theta(\tilde{\Phi}_0^t(\tilde{z}_0)) - \mathcal{X}_\rho \theta(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt,\end{aligned}$$

544 where $\tau^* = \tau^*(I, \theta, s; \rho)$ is a non-degenerate zero of the function (6.6).

545 Since $\mathcal{X}_\rho \theta = 0$ we obtain $\mathcal{S}_2^\theta = 0$.

546 The second integral in (7.7) can be simplified as we did in Section 7.3 to analyze the
547 change in actions, and thus, combining both parts of (7.7) proves the following result:

548 **Corollary 7.5.** *For the perturbation $\mathcal{X}^1 = J\nabla H_1 + \mathcal{X}_\rho$ given in (5.9),*

$$\begin{aligned}\theta_\varepsilon^+ - \theta_\varepsilon^- &= \varepsilon \int_{-\infty}^{\infty} \left(\{\theta, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0)) - \{\theta, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) dt \\ &\quad - \varepsilon \int_{-\infty}^{\infty} \left(\{I, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0)) - \{I, H_1\}(\tilde{\Phi}_0^t(\tilde{z}_0^\pm)) \right) t dt \cdot \left(\frac{\partial^2 h_0}{\partial I^2}(I) \right) \\ &\quad + O(\varepsilon^2) \\ &= \varepsilon \int_{-\infty}^{\infty} \left(\{\theta, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \right. \\ 549 \quad &\quad \left. - \{\theta, H_1\}(0, 0, I, \theta + \omega(I)t, s + t) \right) dt \\ &\quad - \varepsilon \int_{-\infty}^{\infty} \left(\{I, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \right. \\ &\quad \left. - \{I, H_1\}(0, 0, I, \theta + \omega(I)t, s + t) \right) t dt \cdot \left(\frac{\partial^2 h_0}{\partial I^2}(I_0) \right) \\ &\quad + O(\varepsilon^2),\end{aligned}\tag{7.8}$$

550 where $\tau^* = \tau^*(I, \theta, s; \rho)$ is a non-degenerate zero of the function (6.6).

551 In the case when $h_0(I) = \frac{I^2}{2}$ and H_1 is as in (3.10) or (3.11), we have $\frac{\partial^2 h_0}{\partial I^2} = 1$, $\{\theta, H_1\} =$
552 0 , and $\{I, H_1\} = -\frac{\partial H_1}{\partial \theta} = a_{10}f(q)\sin\theta$, where $f(q) = \cos q - 1$ or $f(q) = \cos q$, so

$$553 \quad \theta_\varepsilon^+ - \theta_\varepsilon^- = -\varepsilon a_{10} \int_{-\infty}^{\infty} \left(\cos(q_0(\tau^* + t)) - 1 \right) \sin(\theta + \omega(I)t) t dt + O(\varepsilon^2).$$

554 **Remark 7.6.** We remark that both components $\mathcal{S}^I, \mathcal{S}^\theta$ of the vector field generating the
555 scattering map up to $O(\varepsilon^2)$ only depends on the dissipation \mathcal{X}_λ and, therefore, on the param-
556 eter ρ , through the value $\tau^* = \tau^*(I, \theta, s; \rho)$. In fact, in the next section we will see that the

vector field generating the scattering map is a Hamiltonian vector field in the variables (I, θ) up to $O(\varepsilon^2)$, even though the system (3.8) is not symplectic but conformally symplectic. We will show that the scattering map is symplectic in the variables (I, θ) up to $O(\varepsilon^2)$. Moreover, in the case when H_1 is as in (3.10) or (3.11), we will provide an explicit formula for the Hamiltonian vector field that generates the scattering map up to $O(\varepsilon^2)$.

7.5. Symplecticity of the scattering map up to $O(\varepsilon^2)$. In the case that the perturbation is Hamiltonian (which in our case corresponds to $\rho = 0$), it was proven in [DdlLS08] that the scattering map is symplectic and is given by

$$(7.10) \quad \tilde{\sigma}_\varepsilon(I, \theta, s) = \tilde{\sigma}_0(I, \theta, s) + \varepsilon \left(\frac{\partial L^*}{\partial \theta}(I, \theta, s), -\frac{\partial L^*}{\partial I}(I, \theta, s), s \right) + O(\varepsilon^2)$$

for some function (Melnikov potential) L^* which depends on the effect of the Hamiltonian perturbation on the homoclinic orbits of the unperturbed system. More precisely, let

$$(7.11) \quad \mathcal{L}(I, \theta, s) = - \int_{-\infty}^{+\infty} (H_1(p_0(t), q_0(t), I, \theta + \omega(I)t, s + t) - H_1(0, 0, I, \theta + \omega(I)t, s + t)) dt$$

where $\omega(I) = \frac{\partial h_0}{\partial I}(I)$. Let $\tau^* = \tau^*(I, \theta, s)$ be a non-degenerate critical point of the function

$$\tau \mapsto \mathcal{L}(I, \theta - \omega(I)\tau, s - \tau)$$

Then the function L^* referred to in (7.10) is defined by

$$(7.12) \quad L^*(I, \theta, s) = \mathcal{L}(I, \theta - \omega(I)\tau^*, s - \tau^*).$$

An auxiliary function that will be referred to later is the reduced Melnikov potential defined by

$$(7.13) \quad \mathcal{L}^*(I, \bar{\theta}) = L^*(I, \bar{\theta}, 0) \text{ for } \bar{\theta} = \theta - \omega(I)s.$$

In our case the perturbation is not Hamiltonian, but we will see that, nevertheless, the scattering map is symplectic up to $O(\varepsilon^2)$, and is given by

$$(7.14) \quad \begin{aligned} \mathcal{S}^I &= \frac{\partial L_\rho^*}{\partial \theta} \\ \mathcal{S}^\theta &= -\frac{\partial L_\rho^*}{\partial I}, \end{aligned}$$

for some function L_ρ^* that depends on the effect of the Hamiltonian perturbation on the homoclinic orbits of the unperturbed system and also on the dissipation. Our computation is similar to [DdlLS08].

583 **Proposition 7.7.** *The vector field \mathcal{S} generating the scattering map $\tilde{\sigma}_\varepsilon$ up to $O(\varepsilon^2)$ is of the*
 584 *form*

$$585 \quad (7.15) \quad \mathcal{S}(I, \theta, s) = (-J\nabla_{(I, \theta)} L_\rho^*(I, \theta, s), s)$$

586 *for the function $L_\rho^* : V \subset \tilde{\Lambda}_0 \rightarrow \mathbb{R}$ defined below. Let*

$$587 \quad (7.16) \quad \mathcal{L}(I, \theta, s) = - \int_{-\infty}^{+\infty} (H_1(p_0(t), q_0(t), I, \theta + \omega(I)t, s + t) \\ - H_1(0, 0, I, \theta + \omega(I)t, s + t)) dt$$

588 *where $\omega(I) = \frac{\partial h_0}{\partial I}(I)$.*

589 *Let*

$$590 \quad (7.17) \quad A = \int_{-\infty}^{+\infty} p_0^2(t) dt.$$

591 *Let $\tau^* = \tau^*(I, \theta, s; \rho)$ be a non-degenerate critical point of the function*

$$593 \quad \tau \mapsto \mathcal{L}(I, \theta - \omega(I)\tau, s - \tau) + \rho(s - \tau)A.$$

594 *Let L^* be defined by*

$$595 \quad L^*(I, \theta, s) = \mathcal{L}(I, \theta - \omega(I)\tau^*, s - \tau^*).$$

596 *Then the function L_ρ^* is defined by*

$$597 \quad L_\rho^*(I, \theta, s) = L^*(I, \theta, s) + \rho(s - \tau^*)A.$$

598 **Proof.** We claim that

$$599 \quad (7.18) \quad \begin{aligned} \mathcal{S}^I &= \frac{\partial L_\rho^*}{\partial \theta} = \frac{\partial L^*}{\partial \theta} - \rho \frac{\partial \tau^*}{\partial \theta} A \\ \mathcal{S}^\theta &= - \frac{\partial L_\rho^*}{\partial I} = - \frac{\partial L^*}{\partial I} + \rho \frac{\partial \tau^*}{\partial I} A. \end{aligned}$$

600 The first observation is that the non-degenerate zeroes of the function (6.6) are the non-
 602 degenerate critical points of the function

$$603 \quad (7.19) \quad \tau \in \mathbb{R} \mapsto \mathcal{L}(I, \theta - \omega(I)\tau, s - \tau) + \rho(s - \tau)A,$$

604 where \mathcal{L} is given by (7.16) and A is given by (7.17). To see this, first note that by a change
 605 of variables $t - \tau \mapsto t$, we can express $\mathcal{L}(I, \theta - \omega(I)\tau, s - \tau)$ as

$$606 \quad (7.20) \quad \mathcal{L}(I, \theta - \omega(I)\tau, s - \tau) = - \int_{-\infty}^{+\infty} (H_1(p_0(\tau + t), q_0(\tau + t), I, \theta + \omega(I)t, s + t) \\ - H_1(0, 0, I, \theta + \omega(I)t, s + t)) dt$$

Differentiating (7.19) with respect to τ we obtain

$$\int_{-\infty}^{+\infty} (\{h_1, H_1\}(p_0(\tau+t), q_0(\tau+t), I, \theta + \omega(I)t, s+t) - \{h_1, H_1\}(0, 0, I, \theta + \omega(I)t, s+t)) dt - \rho A,$$

so the non-degenerate zeroes of this function are the non-degenerate critical points of (7.19). If $\tau^* = \tau^*(I, \theta, s; \rho)$ is a non-degenerate critical point of the function (7.19), then by the chain rule it follows that

$$\begin{aligned} 0 &= \frac{d}{d\tau} [\mathcal{L}(I, \theta - \omega(I)\tau, s - \tau) + \rho(s - \tau)A]_{|\tau=\tau^*} \\ &= -\frac{\partial \mathcal{L}}{\partial \theta}(I, \theta - \omega(I)\tau^*, s - \tau^*)\omega(I) - \frac{\partial \mathcal{L}}{\partial s}(I, \theta - \omega(I)\tau^*, s - \tau^*) - \rho A. \end{aligned} \quad (7.21)$$

To compute $\frac{\partial L_\rho^*}{\partial \theta} = \frac{\partial}{\partial \theta}(L^*(I, \theta, s) + \rho(s - \tau^*)A)$ in (7.14) we use the chain rule and (7.21) to obtain

$$\begin{aligned} \frac{\partial L_\rho^*}{\partial \theta} &= \frac{\partial \mathcal{L}}{\partial \theta}(I, \theta - \omega(I)\tau^*, s - \tau^*) \left(1 - \omega(I)\frac{\partial \tau^*}{\partial \theta}\right) \\ &\quad + \frac{\partial \mathcal{L}}{\partial s}(I, \theta - \omega(I)\tau^*, s - \tau^*) \left(-\frac{\partial \tau^*}{\partial \theta}\right) - \rho \frac{\partial \tau^*}{\partial \theta} A \\ &= \frac{\partial \mathcal{L}}{\partial \theta}(I, \theta - \omega(I)\tau^*, s - \tau^*). \end{aligned} \quad (7.22)$$

Applying the latter formula to (7.16), using $\frac{\partial H_1}{\partial \theta} = -\{I, H_1\}$, and making the change of variable $t - \tau^* \mapsto t$ we obtain

$$\begin{aligned} \frac{\partial L_\rho^*}{\partial \theta} &= \int_{-\infty}^{+\infty} (\{I, H_1\}(p_0(t), q_0(t), I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*) \\ &\quad - \{I, H_1\}(0, 0, I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*)) dt \\ &= \int_{-\infty}^{+\infty} (\{I, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \\ &\quad - \{I, H_1\}(0, 0, I, \theta + \omega(I)t, s + t)) dt. \end{aligned} \quad (7.23)$$

This integral is the same as the integral (7.4) that appears in the formula for the change of action by the scattering map up to $O(\varepsilon^2)$. Therefore, we conclude that:

$$I_\varepsilon^+ - I_\varepsilon^- = \varepsilon \frac{\partial L^*}{\partial \theta}(I, \theta, s) + O(\varepsilon^2)$$

To compute $-\frac{\partial L_\rho^*}{\partial I} = -\frac{\partial}{\partial I}(L^*(I, \theta, s) + \rho(s - \tau^*)A)$ in (7.14), we use the chain rule and

622 (7.21) to obtain

$$\begin{aligned}
 -\frac{\partial L_\rho^*}{\partial I} &= -\frac{\partial \mathcal{L}}{\partial I}(I, \theta - \omega(I)\tau^*, s - \tau^*) \\
 &\quad - \frac{\partial \mathcal{L}}{\partial \theta}(I, \theta - \omega(I)\tau^*, s - \tau^*) \left(-\frac{\partial \omega}{\partial I}\tau^* - \omega(I)\frac{\partial \tau^*}{\partial I} \right) \\
 &\quad - \frac{\partial \mathcal{L}}{\partial s}(I, \theta - \omega(I)\tau^*, s - \tau^*) \left(-\frac{\partial \tau^*}{\partial I} \right) \\
 (7.24) \quad &\quad + \rho \frac{\partial \tau^*}{\partial I} A \\
 &= -\frac{\partial \mathcal{L}}{\partial I}(I, \theta - \omega(I)\tau^*, s - \tau^*) \\
 &\quad + \frac{\partial \mathcal{L}}{\partial \theta}(I, \theta - \omega(I)\tau^*, s - \tau^*) \frac{\partial \omega}{\partial I}\tau^*.
 \end{aligned}$$

624

625 We express the two terms in (7.24) as integrals

$$\begin{aligned}
 (7.25) \quad & -\frac{\partial \mathcal{L}}{\partial I}(I, \theta - \omega(I)\tau^*, s - \tau^*) = \int_{-\infty}^{+\infty} (\{\theta, H_1\}(p_0(t), q_0(t), I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*) \\
 & \quad - \{\theta, H_1\}(0, 0, I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*)) dt \\
 626 \quad & - \int_{-\infty}^{+\infty} (\{I, H_1\}(p_0(t), q_0(t), I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*) \\
 & \quad - \{I, H_1\}(0, 0, I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*)) \left(\frac{\partial \omega}{\partial I}(I)t \right) dt
 \end{aligned}$$

627

$$\begin{aligned}
 (7.26) \quad & \frac{\partial \mathcal{L}}{\partial \theta}(I, \theta - \omega(I)\tau^*, s - \tau^*) = \int_{-\infty}^{+\infty} (\{I, H_1\}(p_0(t), q_0(t), I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*) \\
 628 \quad & \quad - \{I, H_1\}(0, 0, I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*)) dt
 \end{aligned}$$

629 Above we used that $\frac{\partial H_1}{\partial \theta} = -\{I, H_1\}$ and $\frac{\partial H_1}{\partial I} = \{\theta, H_1\}$.

630 Combining (7.25) and (7.26) in (7.24) we obtain

$$\begin{aligned}
 (7.27) \quad & -\frac{\partial L_\rho^*}{\partial I}(I, \theta, s) = \int_{-\infty}^{+\infty} (\{\theta, H_1\}(p_0(t), q_0(t), I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*) \\
 & \quad - \{\theta, H_1\}(0, 0, I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*)) dt \\
 631 \quad & - \int_{-\infty}^{+\infty} (\{I, H_1\}(p_0(t), q_0(t), I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*) \\
 & \quad - \{I, H_1\}(0, 0, I, \theta + \omega(I)t - \omega(I)\tau^*, s + t - \tau^*)) \left(\frac{\partial \omega}{\partial I}(I)(t - \tau^*) \right) dt
 \end{aligned}$$

632

633 Making the change of variable $t - \tau^* \mapsto t$ and writing $\frac{\partial \omega}{\partial I}(I) = \frac{\partial^2 h_0}{\partial I^2}(I)$ we obtain

$$\begin{aligned}
-\frac{\partial L^*}{\partial I}(I, \theta, s) &= \int_{-\infty}^{+\infty} (\{\theta, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \\
&\quad - \{\theta, H_1\}(0, 0, I, \theta + \omega(I)t, s + t)) dt \\
(7.28) \quad &- \int_{-\infty}^{+\infty} (\{I, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \\
&\quad - \{I, H_1\}(0, 0, I, \theta + \omega(I)t, s + t)) \left(\frac{\partial^2 h_0}{\partial I^2}(I)t \right) dt
\end{aligned}$$

635

636 Since the integrals in (7.28) are computed in terms of the effect of the perturbation on orbits
 637 of the unperturbed system, we have that I in is constant and equal to $I = I(\tilde{z}_0) = I(\tilde{z}_0^\pm)$, and
 638 therefore $\left(\frac{\partial^2 h_0}{\partial I^2}(I) \right)$ can be taken outside of the second integral obtaining:

$$\begin{aligned}
-\frac{\partial L^*}{\partial I}(I, \theta, s) &= \int_{-\infty}^{+\infty} (\{\theta, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \\
&\quad - \{\theta, H_1\}(0, 0, I, \theta + \omega(I)t, s + t)) dt \\
(7.29) \quad &- \int_{-\infty}^{+\infty} (\{I, H_1\}(p_0(\tau^* + t), q_0(\tau^* + t), I, \theta + \omega(I)t, s + t) \\
&\quad - \{I, H_1\}(0, 0, I, \theta + \omega(I)t, s + t)) t dt \cdot \left(\frac{\partial^2 h_0}{\partial I^2}(I) \right)
\end{aligned}$$

640

641 This integral is the same as the integral (7.8) that appears in the formula for the change of
 642 angle by the scattering map.

643 Therefore, we conclude that:

$$\theta_\varepsilon^+ - \theta_\varepsilon^- = -\varepsilon \frac{\partial L^*}{\partial I}(I, \theta, s) + O(\varepsilon^2)$$

645

646 Consider the mapping

$$L_\rho^*(I, \theta, s) = \mathcal{L}(I, \theta - \omega(I)\tau^*, s - \tau^*) + \rho(s - \tau^*)A.$$

648 for $\tau^* = \tau^*(I, \theta, s; \rho)$. Since τ^* is a critical point for

$$\tau \mapsto \mathcal{L}(I, \theta - \omega(I)\tau, s - \tau) + \rho(s - \tau)A,$$

650 then for every $t' \in \mathbb{R}$, $\tau^* - t'$ is a critical point for

$$\tau \mapsto \mathcal{L}(I, \theta - \omega(I)(\tau + t'), s - (\tau + t')) + \rho(s - (\tau + t'))A.$$

652 Then, denoting $Z = (I, \theta, s; \rho)$ and $Z' = (I, \theta - \omega(I)t', s - t'; \rho)$, we have

$$\tau^*(Z') = \tau^*(Z) - t'.$$

653

Therefore

$$\begin{aligned}
 L_\rho^*(I, \theta - \omega(I)t', s - t') &= \mathcal{L}(I, \theta - \omega(I)(\tau^*(Z') + t'), s - (\tau^*(Z') + t')) \\
 &\quad + \rho(s - (\tau^*(Z') + t'))A \\
 &= \mathcal{L}(I, \theta - \omega(I)(\tau^*(Z) - t' + t'), s - (\tau^*(Z) - t' + t')) \\
 &\quad + \rho(s - (\tau^*(Z) - t' + t'))A \\
 &= L_\rho^*(I, \theta, s).
 \end{aligned}
 \tag{7.30}$$

Making $t' = s$ in (7.30) we obtain

$$L_\rho^*(I, \theta, s) = L_\rho^*(I, \theta - \omega(I)s, 0).$$

This says that, while the function L_ρ^* nominally depends on three variables (I, θ, s) , in fact it depends on the variable I and the linear combination $\theta - \omega(I)s$, and is therefore a function of two independent variables I and $\bar{\theta} = \theta - \omega(I)s$. Thus, we define the reduced Melnikov potential by:

$$\mathcal{L}_\rho^*(I, \bar{\theta}) = \mathcal{L}_\rho^*(I, \bar{\theta}, 0) = \mathcal{L}(I, \bar{\theta} - \omega(I)\tau^*, -\tau^*) - \rho\tau^*A, \text{ for } \bar{\theta} = \theta - \omega(I)s.$$

The reduced Melnikov potential allows to compute the scattering map associated to the time- 2π map associated to a surface of section $\{s = s^*\}$; more precisely, the trajectories of the scattering map are given by the ε -time of the Hamiltonian $-\mathcal{L}_\rho^*$ up to order $O(\varepsilon^2)$, as we shall see below.

7.6. Growth of action by the scattering map. We reduce the dynamics of the flow $\tilde{\Phi}_\varepsilon^t$ to the dynamics of the Poincaré first return map to the surface of section

$$\Sigma = \{(p, q, I, \theta, s) \mid s = s^*\}$$

for some choice of $s^* \in \mathbb{T}^1$.

The NHIM $\tilde{\Lambda}_\varepsilon$ for $\tilde{\Phi}_\varepsilon^t$ in the extend phase space yields the NHIM Λ_ε for f_ε in Σ . In particular Λ_ε is invariant under f_ε .

The scattering map $\tilde{\sigma}_\varepsilon$, which is defined on the domain $\tilde{U} \subseteq \tilde{\Lambda}_\varepsilon$, yields a scattering map σ_ε defined on the following domain in $\Lambda_\varepsilon = \Lambda_0$:

$$U = \{(I, \bar{\theta}) \mid (I, \theta, s^*) \in \tilde{U} \text{ for } \theta = \bar{\theta} + \omega(I)s^*\},$$

The scattering map σ_ε is given in the variables $(I, \bar{\theta})$ by (see [DS18]):

$$\sigma_\varepsilon(I, \bar{\theta}) = \sigma_0(I, \bar{\theta}) - \varepsilon J \nabla \mathcal{L}_\rho^*(I, \bar{\theta}) + O(\varepsilon^2),$$

where $\sigma_0 = \text{Id}$. In particular, the scattering map σ_ε is symplectic up to $O(\varepsilon^2)$.

By Theorem 3.11 in [GdlMS20], whenever $J \nabla \mathcal{L}_\rho^*(z_0) \neq 0$ for some point $z_0 \in \Lambda_0$, there exists a $O(1)$ -family of solutions $\gamma_z(t)$ of the differential equation

$$\dot{z} = -J \nabla \mathcal{L}_\rho^*(z)$$

for z in a $O(1)$ -neighborhood of $z_0 \in U \subseteq \Lambda_0$, and t in some interval $[T_1(z), T_2(z)] \subset \mathbb{R}$ depending on z , such that for each path γ_z , there is an orbit of the scattering map σ_ε that follows closely that path.

If, in addition, we have that $\mathcal{S}^I(z_0) = \frac{\partial \mathcal{L}_\rho^*}{\partial \theta}(z_0) > 0$, then the family of paths γ_z can be chosen so that the corresponding orbits of the scattering map σ_ε along γ_z have the property that I increases by $O(\varepsilon)$ for each application of σ_ε .

Consequently, letting $z_0 = (I_0, \theta_0)$ there exist $\theta_1 < \theta_0 < \theta_2$, $I_1 < I_0 < I_2$, and a ‘strip’ of the form

$$(7.33) \quad S = \{\gamma_z(t) \mid z = (I_0, \theta) \mid \theta \in [\theta_1, \theta_2], t \in [T_1(z), T_2(z)]\} \subseteq U$$

with $\gamma_z(T_1(z)) = I_1$ and $\gamma_z(T_2(z)) = I_2$, such that the following properties hold: There exist $c > 0$, such that for every $\delta = O(\varepsilon) > 0$ and every path $\gamma_z(t)$ contained in S , there exists an orbit $(z_n)_{n=0, \dots, N}$ of σ_ε and $0 = t_0 < t_1 < \dots < t_N = T$ with $t_i = \varepsilon i$ for all i , such that

$$(7.34) \quad \begin{aligned} z_{i+1} &= \sigma_\varepsilon(z_i), \\ I(z_{i+1}) - I(z_i) &> c\varepsilon, \text{ for } i = 0, \dots, N-1, \\ d(z_i, \gamma_z(t_i)) &< \delta, \text{ for } i = 0, \dots, N. \end{aligned}$$

7.7. Scattering map in the case of vanishing and non-vanishing perturbation. For both the vanishing and non-vanishing perturbations:

$$\begin{aligned} H_1(p, q, I, \theta, s) &= (\cos(q) - 1)(a_{00} + a_{10} \cos(\theta) + a_{01} \cos(s)) \\ H_1(p, q, I, \theta, s) &= \cos(q)(a_{00} + a_{10} \cos(\theta) + a_{01} \cos(s)) \end{aligned}$$

we have the same expression for the Melnikov potential

$$(7.35) \quad \begin{aligned} \mathcal{L}(I, \theta, s) &= - \int_{-\infty}^{+\infty} (\cos(q_0(t)) - 1)(a_{00} + a_{10} \cos(\theta + It) + a_{01} \cos(s + t)) dt \\ &= - \int_{-\infty}^{+\infty} (\cos(\arctan e^{2t}) - 1)(a_{00} + a_{10} \cos(\theta + It) + a_{01} \cos(s + t)) dt. \end{aligned}$$

Above we have used the parametrization (3.4) of the separatrix. Since $\frac{p_0^2}{2} + (\cos q_0(t) - 1) = 0$, in the above integral we can alternatively write $\cos(q_0(t)) - 1 = -\frac{p_0^2}{2} = -\frac{2}{\cosh^2(t)}$. It turns out that (see [DG00, DS17])

$$(7.36) \quad \begin{aligned} \mathcal{L}(I, \theta, s) &= A_{00} + A_{10}(I) \cos(\theta) + A_{01}(I) \cos(s), \text{ where} \\ A_{00} &= 4a_{00}, \quad A_{10} = \frac{2\pi I a_{10}}{\sinh(\frac{\pi I}{2})}, \quad A_{01} = \frac{2\pi a_{01}}{\sinh(\frac{\pi}{2})}. \end{aligned}$$

In [DS17] the reduced Melnikov function \mathcal{L}^* defined by (7.13), which corresponds to the Hamiltonian perturbation only, is computed explicitly. The level curves of \mathcal{L}^* are shown in Fig. 5. One can find explicitly regions of size $O(1)$ in Λ_ε where $\frac{\partial \mathcal{L}^*}{\partial \theta}(I, \bar{\theta}) > c_1 > 0$, for some $c_1 > 0$.




Figure 5. Level sets of \mathcal{L}^* . (Credit A. Delshams and R. Schaefer)

In our case, when the system is also subject to the dissipative perturbation \mathcal{X}_λ , the reduced Melnikov potential \mathcal{L}_ρ^* , given by (7.31), is $O(\rho)$ -close to the reduced Melnikov potential \mathcal{L}^* corresponding to the Hamiltonian perturbation. This implies that, for ρ sufficiently small, there exists a region of $O(1)$ in Λ_ε where $\frac{\partial \mathcal{L}_\rho^*}{\partial \theta}(I, \bar{\theta}) > c_2 > 0$, for some $0 < c_2 < c_1$. In the case when $\rho = \frac{\bar{\rho}}{\log(1/\varepsilon)}$, it follows that there exists $\varepsilon_2 > 0$ such that, for all $0 < \varepsilon < \varepsilon_2$, we have that $\frac{\partial \mathcal{L}_\rho^*}{\partial \theta}(I, \bar{\theta}) > c_2 > 0$ on the aforementioned region. This region can be used to define a strip S as in (7.33), where the scattering map increases the action I by $c\varepsilon$ at each step, as in (7.34), for some $0 < c < c_2$.

8. Proof of Theorem 4.1.

8.1. The case of vanishing perturbation. Choose ω_* such that $I_1 < \omega_* < I_2$, where I_1, I_2 are as in Section 7.6. There is an invariant circle $A_\varepsilon = \{I = \omega_*\}$ in Λ_ε , as defined in Section 5.7, which is a global attractor for f_ε on Λ_ε . The circle A_ε is a NHIM for f_ε restricted to Λ_ε , and has only stable manifold $W_{\Lambda_\varepsilon}^s(A_\varepsilon)$ which is the whole manifold Λ_ε . $W_{\Lambda_\varepsilon}^s(A_\varepsilon)$ is foliated by stable leaves

$$W_{\Lambda_\varepsilon}^s(A_\varepsilon) = \bigcup_{y \in A_\varepsilon} W_{\Lambda_\varepsilon}^s(y) \text{ with } y = (\omega_*, \theta) \in A_\varepsilon.$$

From (5.16) we have that each stable leaf is a slanted line

$$W_{\Lambda_\varepsilon}^s(\omega_*, \theta_0) = \left\{ (I, \theta(I)) \in \Lambda_\varepsilon \mid \theta(I) = \theta_0 - \frac{1}{\lambda}(I - \omega_*) \right\}$$

Since $\frac{1}{\lambda} \gg 1$, the slope of these lines (as a function of θ) is $-\lambda$, so the stable leaves are nearly horizontal lines. See Fig. 6.

For $z \in W_{\Lambda_\varepsilon}^s(y)$, by the equivariance property of the stable fibers we have $f_\varepsilon^k(z) \in W_{\Lambda_\varepsilon}^s(f_\varepsilon^k(y))$, for all $k > 0$. Given some initial point (I_0, θ_0) , let $f_\varepsilon^k(I_0, \theta_0) = (I_k, \theta_k)$. From




Figure 6. The attractor A_ε inside the NHIM Λ_ε

(5.15), we deduce

$$\begin{aligned}
 \theta_k &= \theta_0 + \frac{1}{\lambda}(I_0 - \omega_*)(1 + \dots + e^{-2\pi(k-1)\lambda})(1 - e^{-2\pi\lambda}) + 2\pi k\omega_* \\
 &= \theta_0 + \frac{1}{\lambda}(I_0 - \omega_*)(1 - e^{-2\pi k\lambda}) + 2\pi k\omega_* \\
 &= \theta_0 + (I_0 - \omega_*)2\pi k + 2\pi k\omega_* + O(k^2\lambda) \\
 &= \theta_0 + 2\pi kI_0(1 + O(k\lambda)).
 \end{aligned}
 \tag{8.1}$$

Recall that $\lambda = \varepsilon\rho(\varepsilon) = \varepsilon\frac{\bar{\rho}}{\log(\frac{1}{\varepsilon})}$, hence the relative error term in (8.1) is $O(k\frac{\bar{\rho}\varepsilon}{\log(\frac{1}{\varepsilon})})$, so k iterations of the inner map change the angle coordinate by approximately $2\pi kI_0 \pmod{2\pi}$ provided that $k\frac{\bar{\rho}\varepsilon}{\log(\frac{1}{\varepsilon})} \ll 1$.

Consider the strip S defined in (7.33), where the scattering map is increasing I by $O(\varepsilon)$ at each step. Provided that $|I_1 - I_2|$ is suitably small (but independent of ε), there exists $k_{\max} > 0$ such that, whenever $z \in S$ we have $f_\varepsilon^k(z) \in S$ for some $k \leq k_{\max}$. That is, each point z in the strip returns to the strip in a maximum of k_{\max} iterates. This implies that for a time $T = T_0 \log(1/\varepsilon)$, with $\varepsilon > 0$ small, each point z in the strip returns to the strip for at least $\lfloor \frac{T_0 \log(1/\varepsilon)}{k_{\max}} \rfloor$ times.

One easily obtains from (5.15):

$$I_k = (I_0 - \omega_*)e^{-2\pi k\lambda} + \omega_*.$$

Consequently, if $z = (I_0, \theta_0) \in W_{\Lambda_\varepsilon}^s(y)$, $y = (\omega_*, \theta) \in A_\varepsilon$, is a point at a d_I -distance $d_0 \leq d_{\max}$ from y , where $d_{\max} = \max\{|I_1 - \omega_*|, |I_2 - \omega_*|\}$, then $f_\varepsilon^k(z) \in W_{\Lambda_\varepsilon}^s(f_\varepsilon^k(y))$ is at a d_I -distance at most

$$d_0 \cdot e^{-2\pi\lambda k} = d_0 \cdot e^{-2\pi\frac{\varepsilon}{\log(1/\varepsilon)}\bar{\rho}k}$$

from A_ε after k -iterates. For points with initial I_0 above ω_* the I -coordinate decreases at each iterate, and for points with initial I below ω_* the I -coordinate increases at each iterate. Therefore the loss in I after k -iterates, for points with initial $I_0 > \omega_*$, is

$$I_0 - I_k = d_0 \cdot (1 - e^{-2\pi\lambda k}) = d_0 \cdot (1 - e^{-2\pi\frac{\varepsilon}{\log(1/\varepsilon)}\bar{\rho}k}).$$

Hence the maximum loss in the action coordinate I of a point z after k iterates, where $2\pi k \leq T_0 \log(1/\varepsilon)$, is

$$d_{\max} \cdot (1 - e^{-\varepsilon\bar{\rho}T_0}).$$

On the other hand, for each $z \in S$ we can apply a scattering map σ_ε to z . The effect of the scattering map is an increase in the action coordinate I by $O(\varepsilon)$. We recall that there exists $c > 0$ such that

$$I(\sigma_\varepsilon(z)) - I(z) > c\varepsilon, \text{ for all } z \in S.$$

Thus, starting with a point $z \in S$, applying the scattering map σ_ε to z , and then applying the inner map $(f_\varepsilon)|_{\Lambda_\varepsilon}$ for k iterates with $2\pi k \leq T_0 \log(1/\varepsilon)$, until $(f_\varepsilon)^k(\sigma_\varepsilon(z)) \in S$, we obtain a net growth in I that is at least

$$c\varepsilon - d_{\max} \cdot (1 - e^{-\varepsilon \bar{\rho} T_0}).$$

We want that the net growth is at least $c\varepsilon/2$, that is

$$c\varepsilon - d_{\max} \cdot (1 - e^{-\varepsilon \bar{\rho} T_0}) > \frac{c\varepsilon}{2}.$$

This is equivalent to

$$e^{-\varepsilon \bar{\rho} T_0} > 1 - \frac{c\varepsilon}{2d_{\max}}.$$

Taking the logarithm of both sides we obtain

$$T_0 < \frac{\log\left(1 - \frac{c\varepsilon}{2d_{\max}}\right)}{-\varepsilon \bar{\rho}}.$$

By L'Hopital rule

$$\lim_{\varepsilon \rightarrow 0} \frac{\log\left(1 - \frac{c\varepsilon}{2d_{\max}}\right)}{-\varepsilon \bar{\rho}} = \frac{c}{2d_{\max} \bar{\rho}}.$$

This means that, in order to be able to achieve a growth in I of at least $c\varepsilon/2$ per step, for all ε sufficiently small, we need to choose $\bar{\rho}$ small enough so that

$$(8.2) \quad \bar{\rho} < \frac{c}{2d_{\max} T_0}.$$

With these choices, we obtain orbits of the iterated function system (IFS) generated by $\{f_\varepsilon, \sigma_\varepsilon\}$ of the form $z_{n+1} = f_\varepsilon^{k(n)} \circ \sigma_\varepsilon(z_n)$ with $k(n) = O(\log(1/\varepsilon))$, such that I increases by $c\varepsilon/2$ from z_n to z_{n+1} . In $O(\frac{1}{\varepsilon} \log(\frac{1}{\varepsilon}))$ steps, such orbits increase I by $O(1)$. In Section 8.3 we use these orbits of the IFS to produce diffusing pseudo-orbits as claimed as in Theorem 4.1.

Remark 8.1. When a point $z \in \Lambda_\varepsilon$ is of action coordinate $I_0 < \omega_*$ below that of the attractor A_ε , applying the inner dynamics $(f_\varepsilon)|_{\Lambda_\varepsilon}$ to z moves the point towards the attractor, and hence increases I . Thus, the effect of the inner dynamics and the effect of the scattering map concur towards increasing I . The situation reverses when the action coordinate of z is above that of the attractor, in which case the effect of the scattering map is opposed to the effect of the inner dynamics.




Figure 7. The inner dynamics without dissipation.

779 **8.2. The case of non-vanishing perturbation.** The evolution of the I - and θ -variables
 780 under the inner dynamics on $\tilde{\Lambda}_\varepsilon$ is given by:

$$781 \quad (8.3) \quad \begin{cases} \dot{I}_\lambda(t) = -\lambda(I_\lambda(t) - \omega_*) + \varepsilon a_{10} \sin \theta_\lambda(t) \\ \dot{\theta}_\lambda(t) = I_\lambda(t). \end{cases}$$

782
 783 Denote the general solution of the system (8.3) by $z_\lambda(t) = (I_\lambda(t), \theta_\lambda(t))$.
 784 Setting $\lambda = 0$ yields:

$$785 \quad (8.4) \quad \begin{cases} \dot{I}_0(t) = \varepsilon a_{10} \sin \theta_0(t) \\ \dot{\theta}_0(t) = I_0(t), \end{cases}$$

786 with general solution denoted $z_0(t) = (I_0(t), \theta_0(t))$; see Fig. 7. Note that (8.4) is a Hamiltonian
 787 system with Hamiltonian (energy function)

$$788 \quad (8.5) \quad K(I, \theta) = \frac{I^2}{2} + \varepsilon a_{10}(\cos \theta - 1).$$

789
 790 The solutions $(I_0(t), \theta_0(t))$ of the system (8.4) satisfy

$$791 \quad K(I_0(t), \theta_0(t)) = K(I_0(0), \theta_0(0))$$

792 therefore, the application of the inner dynamics does not change the level sets $\{K = \text{const.}\}$
 793 in the case $\lambda = 0$. On the other hand, the variable I may change by up to $O(\varepsilon^{1/2})$ by one
 794 application of the inner dynamics (8.4). At the same time, the change in I by one application
 795 of the scattering map is $O(\varepsilon) \ll O(\varepsilon^{1/2})$. Therefore, instead of comparing the effects on the
 796 action I by the scattering map and by the inner dynamics, as in Section 8.1, in this case we
 797 want to compare the effects on the energy K by the scattering map and by the inner dynamics.
 798 The next lemma gives the change in the energy K that we obtain after one application of the
 799 scattering map:

Lemma 8.2. Let $\tilde{z}_\varepsilon^+ = \sigma_\varepsilon(\tilde{z}_\varepsilon^-)$, where $\tilde{z}_\varepsilon^+ = (I_\varepsilon^+, \theta_\varepsilon^+, s)$ and $\tilde{z}_\varepsilon^- = (I_\varepsilon^-, \theta_\varepsilon^-, s)$. Then:

$$\begin{aligned} K(\tilde{z}_\varepsilon^+) - K(\tilde{z}_\varepsilon^-) &= I_0(I_\varepsilon^+ - I_\varepsilon^-) + O(\varepsilon^2) \\ &= \varepsilon I_0 \frac{\partial \mathcal{L}^*}{\partial \theta}(I, \theta, s) + O(\varepsilon^2) \\ &= \varepsilon a_{10} I_0 \int_{-\infty}^{\infty} (\cos(q_0(\tau^* + t)) - 1) \sin(\theta + \omega(I)t) dt + O(\varepsilon^2), \end{aligned}$$

where $I_0 = I(\tilde{z}_0)$.

Proof. The proof of this lemma is a similar computation to the ones done to compute the change in actions of the scattering map. In fact, using the formulas for the change in actions and angles, Corollaries 7.3 and 7.5, particularly that $I_\varepsilon^+ - I_\varepsilon^- = O(\varepsilon)$ and $\theta_\varepsilon^+ - \theta_\varepsilon^- = O(\varepsilon)$, we have:

$$\begin{aligned} K(\tilde{z}_\varepsilon^+) - K(\tilde{z}_\varepsilon^-) &= \frac{(I_\varepsilon^+)^2}{2} - \frac{(I_\varepsilon^-)^2}{2} + \varepsilon a_{10} (\cos \theta_\varepsilon^+ - \cos \theta_\varepsilon^-) \\ &= \frac{(I_\varepsilon^+ + I_\varepsilon^-)}{2} (I_\varepsilon^+ - I_\varepsilon^-) + \varepsilon a_{10} (\cos \theta_\varepsilon^+ - \cos \theta_\varepsilon^-) \\ &= (I_0 + O(\varepsilon))(I_\varepsilon^+ - I_\varepsilon^-) + O(\varepsilon^2) \\ &= I_0(I_\varepsilon^+ - I_\varepsilon^-) + O(\varepsilon^2). \end{aligned}$$

Applying Proposition 7.7 yields the desired result. ■

The next lemma compares the actions and angles of solutions of systems (8.3) and (8.4).

Lemma 8.3. Let $(I_\lambda(t), \theta_\lambda(t))$ a solution of system (8.3) with initial condition (I_0, θ_0) and $(I_0(t), \theta_0(t))$ the solution of system (8.4) with the same initial condition. Then, there exists $d' > d_{\max} > 0$ such that, for $|\varepsilon|$ small enough, and for $0 \leq t \leq T_0 \log(\frac{1}{\varepsilon})$ we have:

$$\begin{aligned} |I_\lambda(t) - I_0(t)| &\leq d'(1 - e^{-\lambda t}) \\ |\theta_\lambda(t) - \theta_0(t)| &\leq d' \lambda \frac{t^2}{2}. \end{aligned}$$

Proof. Calling

$$u(t) = I_\lambda(t) - I_0(t), \quad \alpha(t) = \theta_\lambda(t) - \theta_0(t),$$

one can easily see that:

$$\begin{aligned} \dot{u} &= \lambda(I_0(t) - \omega_* - u) + \varepsilon a_{10} (\sin(\theta_0(t) + \alpha) - \sin \theta_0(t)) \\ \dot{\alpha} &= u. \end{aligned}$$

Therefore:

$$\begin{aligned} u(t) &= \int_0^t e^{-\lambda(t-s)} [\lambda(I_0(s) - \omega_*) + \varepsilon a_{10} (\sin(\theta_0(s) + \alpha(s)) - \sin \theta_0(s))] ds \\ \alpha(t) &= \int_0^t u(s) ds. \end{aligned}$$

For the first equation in (8.10) we have used the method of variation of constants.

We will bound u in four steps:

821. First we will use the first equation of (8.10) to obtain a weak bound for u ;
 822. Second, we will use the obtained bound on u in the second equation of (8.10) to obtain a
 824 bound for α ;
 823. Third, we will use the obtained bound on α in the first equation of (8.10) to obtain a sharper
 826 bound for u ;
 824. Finally, using the sharper bound on u in the second equation of (8.10) to obtain a new bound
 828 for α .
 829 We will use that the solution $I_0(t)$ is a bounded function, in fact $|I_0(t) - \omega_*| \leq d_{\max}$, where
 830 we recall $d_{\max} = \max\{|I_1 - \omega_*|, |I_2 - \omega_*|\}$. From the first equation of (8.10) we obtain:

$$\begin{aligned} |u(t)| &\leq \int_0^t e^{-\lambda(t-s)} (\lambda d_{\max} + 2\varepsilon a_{10}) ds \\ &\leq \left(d_{\max} + 2\frac{\varepsilon}{\lambda} a_{10}\right) (1 - e^{-\lambda t}). \end{aligned} \quad (8.11)$$

832 Using this bound in the second equation of (8.10) and that $0 \leq 1 - e^{-\lambda t} \leq \lambda t$, we obtain:

$$\begin{aligned} |\alpha(t)| &\leq \int_0^t \left(d_{\max} + 2\frac{\varepsilon}{\lambda} a_{10}\right) \lambda s ds \\ &= \left(d_{\max} + 2\frac{\varepsilon}{\lambda} a_{10}\right) \lambda \frac{t^2}{2}. \end{aligned} \quad (8.12)$$

834 Finally, we will use the obtained bound on α , and the fact that $|\sin(\theta_0(t) + \alpha) - \sin \theta_0(t)| \leq |\alpha|$
 835 in the first equation of (8.10) to obtain:
 836

$$\begin{aligned} |u(t)| &\leq d_{\max}(1 - e^{-\lambda t}) + \varepsilon a_{10} \int_0^t e^{-\lambda(t-s)} |\alpha(s)| ds \\ &\leq d_{\max}(1 - e^{-\lambda t}) + \varepsilon a_{10} \left(d_{\max} + 2\frac{\varepsilon}{\lambda} a_{10}\right) \frac{\lambda t^2}{2} \int_0^t e^{-\lambda(t-s)} ds \\ &\leq \left[d_{\max} + \varepsilon a_{10} \left(d_{\max} + 2\frac{\varepsilon}{\lambda} a_{10}\right) \frac{t^2}{2}\right] (1 - e^{-\lambda t}). \end{aligned} \quad (8.13)$$

838 Observe that, as $0 \leq t \leq T = T_0 \log\left(\frac{1}{\varepsilon}\right)$, and $\lambda = \varepsilon \frac{\bar{\rho}}{\log\left(\frac{1}{\varepsilon}\right)}$ we have that:

$$\varepsilon a_{10} \left(d_{\max} + 2\frac{\varepsilon}{\lambda} a_{10}\right) \frac{t^2}{2} \leq \varepsilon \log^2\left(\frac{1}{\varepsilon}\right) \left(a_{10} d_{\max} \frac{T_0^2}{2}\right) + \varepsilon \log^3\left(\frac{1}{\varepsilon}\right) \left(\frac{a_{10}^2}{\bar{\rho}} T_0^2\right)$$

840 which is arbitrarily small if ε is small (indeed, $\varepsilon \log^p\left(\frac{1}{\varepsilon}\right) \ll \varepsilon^\nu$ for $p > 0$ and $\nu \in (0, 1)$).
 841 Therefore there exist $d' > d_{\max}$ such that:

$$|u(t)| \leq d'(1 - e^{-\lambda t}) \quad (8.14)$$

843 and consequently we have:

$$|I_\lambda(t) - I_0(t)| \leq d'(1 - e^{-\lambda t})$$

for $0 \leq t \leq T_0 \log\left(\frac{1}{\varepsilon}\right)$. Note that d' can be chosen arbitrarily close to d_{\max} provided that ε is small enough.

Using this new bound in the second equation of (8.10) and that $0 \leq 1 - e^{-\lambda t} \leq \lambda t$, we obtain:

$$(8.15) \quad |\theta_\lambda(t) - \theta_0(t)| \leq d' \lambda \frac{t^2}{2}. \quad \blacksquare$$

The next lemma estimates the change in the energy K by the inner dynamics over time intervals of order $O\left(\log\left(\frac{1}{\varepsilon}\right)\right)$.

Lemma 8.4. *There exists $d'' > 0$, such that for $|\varepsilon|$ small enough, and for $0 \leq t \leq T_0 \log\left(\frac{1}{\varepsilon}\right)$ we have:*

$$(8.16) \quad |K(z_\lambda(t)) - K(z_0(t))| \leq d''(1 - e^{-\lambda t}).$$

Proof. From (8.5) and Lemma 8.3 we obtain

$$(8.17) \quad \begin{aligned} |K(z_\lambda(t)) - K(z_0(t))| &= \left| \frac{1}{2}(I_\lambda^2(t) - I_0^2(t)) + \varepsilon a_{10}(\cos(\theta_\lambda(t)) - \cos(\theta_0(t))) \right| \\ &\leq \left| \frac{1}{2}(I_\lambda(t) - I_0(t))(I_\lambda(t) + I_0(t)) \right| + \varepsilon a_{10}|\theta_\lambda(t) - \theta_0(t)| \\ &\leq \left| \frac{1}{2}(I_\lambda(t) - I_0(t))(2I_0(t) + (I_\lambda(t) - I_0(t))) \right| + \varepsilon a_{10}|\theta_\lambda(t) - \theta_0(t)| \\ &\leq \frac{1}{2}d'(1 - e^{-\lambda t}) \left(2(d_{\max} + \omega_*) + d'(1 - e^{-\lambda t}) \right) + \varepsilon a_{10}d' \lambda \frac{t^2}{2} \\ &\leq d'(d_{\max} + \omega_*)(1 - e^{-\lambda t}) + \frac{(d')^2}{2}(1 - e^{-\lambda t})^2 + \varepsilon a_{10}d' \lambda \frac{t^2}{2}. \end{aligned}$$

Taking into account that $\lambda = \frac{\varepsilon \bar{\rho}}{\log\left(\frac{\varepsilon}{2}\right)}$, for $|\varepsilon|$ sufficiently small we have

$$\begin{aligned} (1 - e^{-\lambda t})^2 &\leq (1 - e^{-\lambda t}), \\ \varepsilon \lambda \frac{t^2}{2} &\leq (1 - e^{-\lambda t}), \quad \text{for } 0 \leq t \leq T_0 \log\left(\frac{1}{\varepsilon}\right). \end{aligned}$$

Therefore, using (8.17) we conclude that there exists $d'' > d'$, such that

$$(8.18) \quad |K(z_\lambda(t)) - K(z_0(t))| \leq d''(1 - e^{-\lambda t}).$$

We note that d'' can be chosen arbitrarily close to $d'(d_{\max} + \omega_*) + \frac{1}{2}(d')^2$ provided that $|\varepsilon|$ is sufficiently small. \blacksquare

We now continue with the proof of Theorem 4.1. By Lemma 8.2, given that $0 < I_1 < I_2$, the effect of the scattering map is an increase in the energy K by $O(\varepsilon)$. Let S be a strip as in (7.33) and $c > 0$ such that

$$(8.19) \quad K(\sigma_\varepsilon(z)) - K(z) > c\varepsilon, \quad \text{for all } z \in S.$$

864 From Lemma 8.4, the maximum loss in K by the inner flow over a time $0 \leq t \leq T_0 \log(\frac{1}{\varepsilon})$ is

$$865 \quad d''(1 - e^{-\varepsilon \bar{\rho} T_0}).$$

866 Switching from the flow to the time- 2π map, it follows that the maximum loss in K by the
867 inner map after k iterates of f_ε , with $2\pi k \leq T_0 \log(1/\varepsilon)$, is also

$$868 \quad d''(1 - e^{-\varepsilon \bar{\rho} T_0}).$$

869 Note that the level sets of K are $O(\varepsilon^{1/2})$ -close to level sets of I . Therefore we can choose
870 $0 < I_1 < I_2$ such that the growth in K by repeated applications of the scattering map
871 corresponds to a change in I from below I_1 to above I_2 . Provided that $|I_1 - I_2|$ is suitably
872 small, there exists k_{\max} such that whenever $z \in S$ we have $f_0^k(z) \in S$ for some $k \leq k_{\max}$. By
873 Lemma 8.8, for $0 \leq t \leq T_0 \log(1/\varepsilon)$,

$$874 \quad |\theta_\lambda(t) - \theta_0(t)| < d' \lambda \frac{t^2}{2} \leq d' \varepsilon \log\left(\frac{1}{\varepsilon}\right) \frac{\bar{\rho} T_0^2}{2},$$

875 so, for ε sufficiently small, $\theta_\lambda(t)$ is $O(\varepsilon^\nu)$ -close to $\theta_0(t)$, for some $\nu \in (0, 1)$. This implies that
876 each point z in the strip returns to the strip in a maximum of k_{\max} iterates of f_ε . Thus,
877 starting with a point $z \in S$, applying the scattering map σ_ε to z , and then applying the inner
878 map $(f_\varepsilon)|_{\Lambda_\varepsilon}$ for k times, where $2\pi k \leq T_0 \log(1/\varepsilon)$, until $(f_\varepsilon)^k(\sigma_\varepsilon(z)) \in S$, we obtain a net
879 growth in K that is at least

$$880 \quad c\varepsilon - d''(1 - e^{-\varepsilon \bar{\rho} T_0}).$$

881 We require that this net growth in K is at least $c\varepsilon/2$, that is

$$882 \quad c\varepsilon - d''(1 - e^{-\varepsilon \bar{\rho} T_0}) > \frac{c}{2}\varepsilon.$$

Similarly to the proof in Section 8.1, in order to be able to achieve a growth in K of at least $c\varepsilon/2$ per step, for all ε sufficiently small, we need to choose $\bar{\rho}$ small enough so that

$$\bar{\rho} < \frac{c}{2d''T_0}.$$

883 We obtain orbits of the iterated function system (IFS) generated by $\{f_\varepsilon, \sigma_\varepsilon\}$, of the form
884 $z_{n+1} = f_\varepsilon^{k(n)} \circ \sigma_\varepsilon(z_n)$ with $k(n) = O(\log(1/\varepsilon))$, such that K increases by $c\varepsilon/2$ from z_n to z_{n+1} .
885 In $O(\frac{1}{\varepsilon} \log(\frac{1}{\varepsilon}))$ steps, such orbits increase K , as well as I , by $O(1)$. By our choice of I_1, I_2 ,
886 these orbits go from below I_1 to above I_2 . In Section 8.3 we use these orbits of the IFS to
887 produce diffusing pseudo-orbits as claimed in Theorem 4.1.

888 **8.3. Existence of diffusing pseudo-orbits.** In Sections 8.1 and 8.2 we obtained diffusing
889 orbits of the iterated function system (IFS) generated by $\{f_\varepsilon, \sigma_\varepsilon\}$ consisting of orbit segments
890 of the form $z_{n+1} = f_\varepsilon^{k(n)} \circ \sigma_\varepsilon(z_n)$ in Λ_ε , $n = 1, \dots, m-1$, where $k(n) = O(\log(1/\varepsilon))$ and
891 $m = O(1/\varepsilon)$, such that $I(z_0) < I_1$ and $I(z_m) > I_2$. We can rearrange these orbits into orbit
892 segments of the form $x_{i+1} = f^{m_i} \circ \sigma_\varepsilon \circ f^{n_i}(x_i)$, with $m_i, n_i = O(\log(1/\varepsilon))$, for $i = 0, \dots, m-1$,
893 such that $I(x_0) < I_1$ and $I(x_m) > I_2$. Each such orbit segment can be approximated up to

$O(\varepsilon)$ by a true orbit of the Poincaré map of the form $y_{\text{end}}^i = f^{k_i}(y_{\text{start}}^i)$, $k_i = O(\log(1/\varepsilon))$, with $d(y_{\text{end}}^i, y_{\text{start}}^{i+1}) < O(\varepsilon)$; see Remark 7.1. The diffusion time is $O(\frac{1}{\varepsilon} \log(\frac{1}{\varepsilon}))$. Finally, each such orbit of f_ε gives rise to an orbit segment $z^i(t)$, $t \in [t_i, t_{i+1}]$, of the flow, satisfying the requirements of Theorem 4.1.

REFERENCES

- [Arn64] V.I. Arnold. Instability of dynamical systems with several degrees of freedom. *Sov. Math. Doklady*, 5:581–585, 1964.
- [BB02] Massimiliano Berti and Philippe Bolle. A functional analysis approach to Arnold diffusion. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 19(4):395–450, 2002.
- [BBB03] Massimiliano Berti, Luca Biasco, and Philippe Bolle. Drift in phase space: a new variational mechanism with optimal diffusion time. *J. Math. Pures Appl. (9)*, 82(6):613–664, 2003.
- [BG05] Keith Burns and Marian Gidea. *Differential geometry and topology: With a view to dynamical systems*. Studies in Advanced Mathematics. Chapman Hall/CRC, Boca Raton, FL, 2005.
- [BKZ16] Patrick Bernard, Vadim Kaloshin, and Ke Zhang. Arnold diffusion in arbitrary degrees of freedom and normally hyperbolic invariant cylinders. *Acta Mathematica*, 217(1):1–79, 2016.
- [BT99] S Bolotin and D Treschev. Unbounded growth of energy in nonautonomous Hamiltonian systems. *Nonlinearity*, 12(2):365, 1999.
- [CC09] Alessandra Celletti and Luigi Chierchia. Quasi-periodic attractors in celestial mechanics. *Archive for rational mechanics and analysis*, 191(2):311–345, 2009.
- [CCdIL13a] Renato C Calleja, Alessandra Celletti, and Rafael De la Llave. A KAM theory for conformally symplectic systems: efficient algorithms and their validation. *Journal of Differential Equations*, 255(5):978–1049, 2013.
- [CCdIL13b] Renato C Calleja, Alessandra Celletti, and Rafael de la Llave. Local behavior near quasi-periodic solutions of conformally symplectic systems. *Journal of Dynamics and Differential Equations*, 25(3):821–841, 2013.
- [CCdIL20] Renato Calleja, Alessandra Celletti, and Rafael de la Llave. KAM theory for some dissipative systems. *arXiv preprint arXiv:2007.08394*, 2020.
- [Cel07] Alessandra Celletti. Weakly dissipative systems in Celestial Mechanics. In *Topics in Gravitational Dynamics*, pages 67–90. Springer, 2007.
- [CG] Maciej J Capiński and Marian Gidea. Arnold diffusion, quantitative estimates, and stochastic behavior in the three-body problem. *Communications on Pure and Applied Mathematics*.
- [CG94] L. Chierchia and G. Gallavotti. Drift and diffusion in phase space. *Ann. Inst. H. Poincaré Phys. Théor.*, 60(1):144, 1994.
- [Chi79] B.V. Chirikov. A universal instability of many-dimensional oscillator systems. *Phys. Rep.*, 52(5):264–379, 1979.
- [CX19] Chong-Qing Cheng and Jinxin Xue. Variational approach to arnold diffusion. *Science China Mathematics*, 62(11):2103–2130, 2019.
- [CY09] Chong-Qing Cheng and Jun Yan. Arnold diffusion in Hamiltonian systems: a priori unstable case. *J. Differential Geom.*, 82(2):229–277, 2009.
- [DdILS00] Amadeu Delshams, Rafael de la Llave, and Tere M. Seara. A geometric approach to the existence of orbits with unbounded energy in generic periodic perturbations by a potential of generic geodesic flows of \mathbf{T}^2 . *Comm. Math. Phys.*, 209(2):353–392, 2000.
- [DdILS06a] Amadeu Delshams, Rafael de la Llave, and Tere M. Seara. A geometric mechanism for diffusion in Hamiltonian systems overcoming the large gap problem: heuristics and rigorous verification on a model. *Mem. Amer. Math. Soc.*, 179(844):viii+141, 2006.
- [DdILS06b] Amadeu Delshams, Rafael de la Llave, and Tere M. Seara. Orbits of unbounded energy in quasi-periodic perturbations of geodesic flows. *Adv. Math.*, 202(1):64–188, 2006.
- [DdILS08] Amadeu Delshams, Rafael de la Llave, and Tere M. Seara. Geometric properties of the scattering map of a normally hyperbolic invariant manifold. *Adv. Math.*, 217(3):1096–1153, 2008.
- [DG00] Amadeu Delshams and Pere Gutiérrez. Splitting potential and the Poincaré-Melnikov method

- for whiskered tori in Hamiltonian systems. *Journal of Nonlinear Science*, 10(4):433–476, 2000.
- [DH09] Amadeu Delshams and Gemma Huguet. Geography of resonances and Arnold diffusion in a priori unstable Hamiltonian systems. *Nonlinearity*, 22(8):1997, 2009.
- [DLS06] Amadeu Delshams, Rafael de la Llave, and Tere M. Seara. A geometric mechanism for diffusion in Hamiltonian systems overcoming the large gap problem: heuristics and rigorous verification on a model. *Mem. Amer. Math. Soc.*, 179(844):viii+141, 2006.
- [DS17] Amadeu Delshams and Rodrigo G. Schaefer. Arnold diffusion for a complete family of perturbations. *Regular and Chaotic Dynamics*, 22(1):78–108, 2017.
- [DS18] Amadeu Delshams and Rodrigo G. Schaefer. Arnold diffusion for a complete family of perturbations with two independent harmonics. *Discrete & Continuous Dynamical Systems - A*, 38(12):6047, 2018.
- [EHI09] Alper Erturk, J Hoffmann, and Daniel J Inman. A piezomagnetoelastic structure for broadband vibration energy harvesting. *Applied Physics Letters*, 94(25):254102, 2009.
- [Fen74] N. Fenichel. Asymptotic stability with rate conditions. *Indiana Univ. Math. J.*, 23:1109–1137, 1973/74.
- [GdlL17] Marian Gidea and Rafael de la Llave. Perturbations of geodesic flows by recurrent dynamics. *J. Eur. Math. Soc. (JEMS)*, 19(3):905–956, 2017.
- [GdlL18] Marian Gidea and Rafael de la Llave. Global Melnikov theory in Hamiltonian systems with general time-dependent perturbations. *Journal of NonLinear Science*, 28:1657–1707, 2018.
- [GdlLM21] Marian Gidea, Rafael de la Llave, and Maxwell Musser. Global effect of non-conservative perturbations on homoclinic orbits. *Qualitative Theory of Dynamical Systems*, 20(1):1–40, 2021.
- [GdlLM22] Marian Gidea, Rafael de la Llave, and Maxwell Musser. Melnikov method for non-conservative perturbations of the restricted three-body problem. *Celestial Mechanics and Dynamical Astronomy*, 134(1):1–42, 2022.
- [GdlLMS20] Marian Gidea, Rafael de la Llave, and Tere M-Seara. A general mechanism of diffusion in Hamiltonian systems: Qualitative results. *Communications on Pure and Applied Mathematics*, 73(1):150–209, 2020.
- [GM22] Marian Gidea and Jean-Pierre Marco. Diffusing orbits along chains of cylinders. *Discrete and Continuous Dynamical Systems*, 0:–, 2022.
- [GR13] Marian Gidea and Clark Robinson. Diffusion along transition chains of invariant tori and Aubry–Mather sets. *Ergodic Theory and Dynamical Systems*, 33(5):1401–1449, 2013.
- [Gra17] Albert Granados. Invariant manifolds and the parameterization method in coupled energy harvesting piezoelectric oscillators. *Physica D: Nonlinear Phenomena*, 351:14–29, 2017.
- [Gre79] John M Greene. A method for determining a stochastic transition. *Journal of Mathematical Physics*, 20(6):1183–1201, 1979.
- [GT08] Vassili Gelfreich and Dmitry Turaev. Unbounded energy growth in Hamiltonian systems with a slowly varying parameter. *Comm. Math. Phys.*, 283(3):769–794, 2008.
- [GT17a] Vassili Gelfreich and Dmitry Turaev. Arnold diffusion in a priori chaotic symplectic maps. *Communications in Mathematical Physics*, 353(2):507–547, Jul 2017.
- [GT17b] Vassili Gelfreich and Dmitry Turaev. Arnold diffusion in a priori chaotic symplectic maps. *Communications in Mathematical Physics*, 353(2):507–547, 2017.
- [HPS77] M.W. Hirsch, C.C. Pugh, and M. Shub. *Invariant manifolds*, volume 583 of *Lecture Notes in Math*. Springer-Verlag, Berlin, 1977.
- [Joh14] Kim Johannessen. An analytical solution to the equation of motion for the damped nonlinear pendulum. *European Journal of Physics*, 35(3):035014, 2014.
- [KZ15] V. Kaloshin and K. Zhang. Arnold diffusion for smooth convex systems of two and a half degrees of freedom. *Nonlinearity*, 28(8):2699–2720, 2015.
- [KZ20] Vadim Kaloshin and Ke Zhang. *Arnold Diffusion for Smooth Systems of Two and a Half Degrees of Freedom:(AMS-208)*. Princeton University Press, 2020.
- [Mat04] John N. Mather. Arnold diffusion. I. Announcement of results. *J. Math. Sci. (N. Y.)*, 124(5):5275–5289, 2004.
- [Mat12] John N. Mather. Arnold diffusion by variational methods. In *Essays in mathematics and its*

- 998 *applications*, pages 271–285. Springer, Heidelberg, 2012.
- 999 [MH79] FC Moon and Philip J Holmes. A magnetoelastic strange attractor. *Journal of Sound and*
1000 *Vibration*, 65(2):275–296, 1979.
- 1001 [MNF87] Andrea Milani, Anna Maria Nobili, and Paolo Farinella. *Non-gravitational perturbations and*
1002 *satellite geodesy*. CRC Press, 1987.
- 1003 [Pif06] G. N. Piftankin. Diffusion speed in the Mather problem. *Dokl. Akad. Nauk*, 408(6):736–737,
1004 2006.
- 1005 [RR17] Clodoaldo Ragazzo and LS Ruiz. Viscoelastic tides: models for use in Celestial Mechanics.
1006 *Celestial Mechanics and Dynamical Astronomy*, 128(1):19–59, 2017.
- 1007 [Tre02] D. Treschev. Evolution of slow variables in near-integrable Hamiltonian systems. In *Progress in*
1008 *nonlinear science, Vol. 1 (Nizhny Novgorod, 2001)*, pages 166–169. RAS, Inst. Appl. Phys.,
1009 Nizhniĭ Novgorod, 2002.
- 1010 [Tre04] D. Treschev. Evolution of slow variables in a priori unstable Hamiltonian systems. *Nonlinearity*,
1011 17(5):1803–1841, 2004.
- 1012 [Tre12] D. Treschev. Arnold diffusion far from strong resonances in multidimensional *a priori* unstable
1013 Hamiltonian systems. *Nonlinearity*, 25(9):2717–2757, 2012.
- 1014 [Zgl09] Piotr Zgliczyński. Covering relations, cone conditions and the stable manifold theorem. *J.*
1015 *Differential Equations*, 246(5):1774–1819, 2009.