

# Addendum to “Commensurations and Subgroups of Finite Index of Thompson’s Group $F$ ”

JOSÉ BURILLO  
SEAN CLEARY  
CLAAS E. RÖVER

It is shown that the abstract commensurator of  $F$  is composed of four building blocks. Two isomorphism types of simple groups, the multiplicative group of the positive rationals and a cyclic group of order two. The main result establishes the simplicity of a certain group of piecewise linear homeomorphisms of the real line.

[20E34](#), [20E32](#); [20F65](#)

The purpose of this note is to extend earlier work [2], where we described the commensurator group of Thompson’s group  $F$ . We prove that an interesting subgroup of  $\text{Com}(F)$  is simple and describe the algebraic structure of  $\text{Com}(F)$  in terms of short exact sequence of simple groups and the multiplicative group of the positive rationals. For all the details and notation, see the paper [2].

## 1 The group of eventually periodic maps

Previously [2] we described the commensurator group of  $F$  as the group of the eventually integrally periodically affine maps in  $P$ , which is defined in Section 1 of [2]. These elements may preserve or reverse the orientation of the real line. We also showed that the index-two subgroup  $\text{Com}^+(F)$  of orientation-preserving maps fits into the short exact sequence

$$1 \longrightarrow K \longrightarrow \text{Com}^+(F) \longrightarrow \mathbb{Q}^* \times \mathbb{Q}^* \longrightarrow 1$$

whose kernel  $K$  is exactly those elements  $f$  of  $P_+$  for which there exists  $M > 0$ , and two positive integers  $p, p'$  such that

$$\begin{aligned} f(t + p) &= f(t) + p \text{ for } t \geq M \text{ and} \\ f(t + p') &= f(t) + p' \text{ for } t \leq -M. \end{aligned}$$

Now we can associate to each element  $f \in K$  two integrally periodically affine maps  $f_+$  and  $f_-$ , which coincide with  $f$  near  $\infty$  and  $-\infty$ , respectively. This property leads to the following definitions.

For  $p \in \mathbb{N}$ , we denote by  $H_p$  the subgroup of  $P_+$  of  $p$ -periodically affine maps, that is

$$H_p = \{f \in P_+ \mid f(t+p) = f(t) + p \text{ for all } t \in \mathbb{R}\}.$$

Clearly, if  $p|q$ , then  $H_p \subset H_q$ , whence we define the subgroup  $H$  as a direct limit under inclusion by

$$H = \bigcup_{p=1}^{\infty} H_p.$$

The maps  $f_+$  and  $f_-$  now give rise to a homomorphism

$$\rho: K \longrightarrow H \times H,$$

given by  $\rho(f) = (f_-, f_+)$ . The kernel consists of the eventually trivial elements, and therefore equals  $F'$ , the commutator subgroup of  $F$  (see [1] or [2]). In other words, we get the short exact sequence

$$1 \longrightarrow F' \longrightarrow K \longrightarrow H \times H \longrightarrow 1.$$

Brin [1] showed that  $\text{Aut}^+(F) = \rho^{-1}(H_1 \times H_1)$  and established the short exact sequence

$$1 \longrightarrow F \longrightarrow \text{Aut}^+(F) \longrightarrow T \times T \longrightarrow 1,$$

where  $T$  is Thompson's group  $T$  (see [3]). Since we clearly have a map  $H_1 \rightarrow T$ , due to the fact that a map which is 1-periodically affine can be viewed as a map on the circle  $S^1$  given by  $\mathbb{R}/\mathbb{Z}$ , an alternative version of this sequence is

$$1 \longrightarrow F' \longrightarrow \text{Aut}^+(F) \longrightarrow H_1 \times H_1 \longrightarrow 1.$$

These two sequences are related by the short exact sequence

$$1 \longrightarrow A_1 \longrightarrow H_1 \longrightarrow T \longrightarrow 1,$$

whose kernel  $A_1$  is the maps  $t \mapsto t + k$  for integers  $k$ . Clearly  $A_1$  is isomorphic to  $\mathbb{Z}$ .

It is straightforward to verify that any element  $\alpha$  of  $\text{Com}^+(F)$  which satisfies  $\alpha(t+1) = \alpha(t) + p$  for all  $t \in \mathbb{R}$  conjugates  $H_1$  to  $H_p$  and  $A_1$  to  $A_p$ , the group of maps of the form  $t \mapsto t + kp$  with  $k \in \mathbb{Z}$ . So we clearly have a short exact sequence

$$1 \longrightarrow A_p \longrightarrow H_p \longrightarrow T \longrightarrow 1.$$

We note that this extension is, in fact, central, and that one may view this copy of  $T$  as acting on the circle of length  $p$  given by  $\mathbb{R}/p\mathbb{Z}$ . We summarise this discussion as follows.

**Theorem 1** *The structure of the group  $\text{Com}(F)$  and its index-two subgroup  $\text{Com}^+(F)$  is given by the following short exact sequences and equalities.*

$$1 \longrightarrow \text{Com}^+(F) \longrightarrow \text{Com}(F) \longrightarrow C_2 \longrightarrow 1$$

$$1 \longrightarrow K \longrightarrow \text{Com}^+(F) \longrightarrow \mathbb{Q}^* \times \mathbb{Q}^* \longrightarrow 1$$

$$1 \longrightarrow F' \longrightarrow K \longrightarrow H \times H \longrightarrow 1, \quad H = \bigcup_{p=1}^{\infty} H_p$$

$$1 \longrightarrow A_p \longrightarrow H_p \longrightarrow T \longrightarrow 1$$

$$A_p \cong \mathbb{Z} \text{ is central in } H_p$$

## 2 Simplicity of the group $H$

Here we exploit the well known fact that  $T$  is simple (eg. [3]) to prove our main result.

**Theorem 2** *The group  $H = \{f \in P_+ \mid f(t+p) = f(t)+p \text{ for some } p \in \mathbb{N}\}$  is simple.*

Note that for  $p, q \in \mathbb{N}$  with  $p|q$ , we have  $H_p \subset H_q$  and  $A_p \supset A_q$ . So the theorem says that in the union  $H$  the groups  $A_p$  cease to be normal. This is due to the following.

**Lemma 3** *A normal subgroup of  $H_p$  is either  $H_p$  or it is contained in  $A_p$ .*

**Proof** In the light of the isomorphism between  $H_p$  and  $H_1$  which carries  $A_p$  to  $A_1$ , it suffices to consider the case  $p = 1$ . Let  $N$  be a normal subgroup of  $H_1$  and consider its image in  $T$ . Since  $T$  is simple, the image of  $N$  is either  $\{1\}$  or the whole  $T$ . If the image is  $\{1\}$ , then  $N \subset A_1$ . So we assume that the image is  $T$ , which yields the exact sequence

$$1 \longrightarrow B \longrightarrow N \longrightarrow T \longrightarrow 1$$

with kernel  $B = N \cap A_1 \subset A_1$ . It follows that  $B = A_r$  for some  $r$ , and we find that

$$H_1/N \cong A_1/A_r \cong \mathbb{Z}/r\mathbb{Z}.$$

In particular  $H_1/N$  is abelian. The proof will be complete once we show that  $H_1$  is equal to its commutator subgroup, because then  $N = H_1$ . On order to establish this, we recall from [3] that  $T$  is generated by three elements  $x_0, x_1$  and  $c$  subject to the relators

$$[x_0x_1^{-1}, x_0^{-1}x_1x_0], \quad [x_0x_1^{-1}, x_0^{-2}x_1x_0^2], \quad x_1x_0^{-1}cx_1c^{-1}, \\ (x_0^{-1}cx_1)^2x_0^{-1}c^{-1}, \quad x_1x_0^{-2}cx_1^2x_0^{-1}x_1^{-1}x_0x_1^{-1}c^{-1}x_0 \quad \text{and} \quad c^3.$$

This easily gives rise to a finite presentation for  $H_1$  with three generators  $x_0$ ,  $x_1$  and  $c$  subject to the same relators, except for  $c^3$  which has to be replaced by the two relators  $[c^3, x_0]$  and  $[c^3, x_1]$ . Here  $x_0$ ,  $x_1$  and  $c$  are the preimages of the corresponding generators for  $T$ , as defined in [3], with  $x_0(0) = x_1(0) = 0$  and  $c(0) = -1/4$ ; composition is then to be read from right to left, as in [3]. In this case  $c^3$  is the map  $t \mapsto t - 1$  which generates  $A_1$ . Modulo the commutator subgroup of  $H_1$ , the third, fourth and fifth relators yield the relators  $x_0^{-1}x_1^2$ ,  $x_0^{-3}x_1^2c$  and  $x_0^{-1}x_1$ , respectively, which in turn imply  $x_0 = x_1 = c = 1$ . This proves that  $[H_1, H_1] = H_1$ .  $\square$

**Proof of Theorem 2.** Let  $N$  be a non-trivial normal subgroup of  $H$ . According to the lemma, for each  $p$ , we have that  $N \cap H_p$  is either  $H_p$  or it is contained in  $A_p$ .

We claim that if  $N \cap H_p = H_p$  for some  $p$ , then this happens for all  $p \in \mathbb{N}$ . Take  $q \in \mathbb{N}$ . Then  $N \cap H_{pq}$  is a normal subgroup of  $H_{pq}$ , and

$$N \cap H_{pq} \supset N \cap H_p = H_p \supsetneq A_p \supset A_{pq},$$

which shows that  $N \cap H_{pq} = H_{pq}$ , by the lemma. Thus, in this case  $N$  contains all  $H_q$ , and hence  $N = H$ .

The only case left now is that  $N \cap H_p \subset A_p$  for all  $p$ . Since  $N$  is non-trivial and the  $A_p$  are infinite cyclic, there exists a  $p$  with  $N \cap H_p = A_{rp}$  for some  $r \geq 1$ . But then

$$A_{2pr} \supset N \cap H_{2pr} \supset N \cap H_p = A_{rp} \supsetneq A_{2pr}$$

which is a contradiction. Thus the only normal subgroups of  $H$  are  $H$  and the identity as claimed.  $\square$

We would like to thank the de Brún Centre at NUI Galway for its generous support. The first author acknowledges support from MEC grant MTM2011-25955 and the second author is grateful for funding from NSF #0811002.

## References

- [1] Matthew G. Brin. The chameleon groups of Richard J. Thompson: automorphisms and dynamics. *Inst. Hautes Études Sci. Publ. Math.*, 84:5–33 (1997), 1996.

- [2] José Burillo, Sean Cleary and Claas E. Röver. Commensurations and finite-index subgroups of Thompson’s group  $F$ . *Geom. Topol.*, 12:1701–1709, 2008.
- [3] J. W. Cannon, W. J. Floyd, and W. R. Parry. Introductory notes on Richard Thompson’s groups. *Enseign. Math. (2)*, 42(3-4):215–256, 1996.

*Departament de Matemàtica Aplicada IV, Universitat Politècnica de Catalunya, Escola Politècnica Superior de Castelldefels, Esteve Terrades 5, 08860 Castelldefels (Barcelona), Spain*

*Department of Mathematics R8133, The City College of New York, Convent Ave & 138th, New York, NY 10031*

*School of Mathematics, Statistics and Applied Mathematics, National University of Ireland Galway, University Road, Galway, Ireland*

[burillo@mat.upc.es](mailto:burillo@mat.upc.es), [cleary@sci.ccnycunyu.edu](mailto:cleary@sci.ccnycunyu.edu), [claas.roever@nuigalway.ie](mailto:claas.roever@nuigalway.ie)

<http://www-ma4.upc.edu/~burillo/>, <http://www.sci.ccnycunyu.edu/~cleary/>,

<http://www.maths.nuigalway.ie/~chew/>